



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07K 14/00	A2	(11) International Publication Number: WO 00/46240 (43) International Publication Date: 10 August 2000 (10.08.00)
(21) International Application Number: PCT/US00/01871 (22) International Filing Date: 27 January 2000 (27.01.00) (30) Priority Data: 09/244,448 3 February 1999 (03.02.99) US 09/264,527 8 March 1999 (08.03.99) US (63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Applications US 09/244,448 (CIP) Filed on 3 February 1999 (03.02.99) US 09/264,527 (CIP) Filed on 8 March 1999 (08.03.99) (71) Applicant (for all designated States except US): AMGEN INC. [US/US]; One Amgen Center Drive, Thousand Oaks, CA 91320-1799 (US). (72) Inventor; and (75) Inventor/Applicant (for US only): YOSHINAGA, Steven, Kiyoshi [US/US]; 644 Lynwood Street, Thousand Oaks, CA 91360 (US).		(74) Agents: ODRE, Steven, M. et al.; Amgen, Inc., One Amgen Center Drive, Thousand Oaks, CA 91320-1799 (US). (81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: NOVEL POLYPEPTIDES INVOLVED IN IMMUNE RESPONSE (57) Abstract <p>Novel polypeptides which comprise a receptor-ligand pair involved in T-cell activation are disclosed. Nucleic acid molecules encoding said polypeptides, and vectors and host cells for expressing same are also disclosed. The polypeptides, or agonists and antagonists thereof, are used to treat T-cell mediated disorders.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

- 1 -

NOVEL POLYPEPTIDES INVOLVED IN IMMUNE RESPONSE

Field of the Invention

The present invention relates to polypeptides
5 that are involved in T-lymphocyte activation.
Specifically, the invention relates to T-lymphocyte
costimulatory polypeptides, the nucleic acids encoding
the polypeptides, expression vectors and host cells for
production of the polypeptides, and compositions and
10 methods for the treatment of diseases related to
immunosuppression and immune activation.

Background of the invention

For the generation of a proper T-lymphocyte
15 (T-cell) immune response, two signals must be provided
to the T-cell by antigen presenting cells (APC). First,
antigen must be presented to the T-cell receptor (TCR)
via a major histocompatibility complex (MHC), in an
event that determines specificity. Second, an antigen-
20 independent, costimulatory signal must be delivered by
engagement of members of the B7 family on the APC with
the CD28 protein on T-cells. A productive immune
response leads to proliferation, differentiation,
clonal expansion, and effect or function. In the
25 absence of the second, costimulatory signal, T-cells
undergo a state of long-lasting antigen-specific
unresponsiveness, termed anergy.

T-cells initiate the immune response, mediate
antigen-specific effector functions, and regulate the
30 activity of other leukocytes by secreting cytokines.
The T-cell receptor (TCR) distinguishes the T-cell from
other lymphocytes and can bind antigen only when it is
presented by the APC within the context of a MHC. The
functional activity of a particular T-cell can be
35 correlated with the expression of membrane antigens,
such as CD4 and CD8. For instance, CD4+ T-cells

- 2 -

generally function as T helper cells (T_H) and are MHC class II restricted, whereas CD8+ cells generally function as cytotoxic T-cells (T_C) and are MHC class I restricted.

5 Potent T-cell costimulatory polypeptides which has been previously identified include polypeptides termed B7.1 (Freeman et al. J. Immunology 143, 2714-2722 (1989), Freeman et al. Jour. Expt. Med. 174, 625-31 (1991)) and B7.2 (Freeman et al. Science 262, 909-911 (1993), and Freeman et al. Jour. Expt. Med. 178, 2185-2192 (1993)), (or CD80 and CD86, respectively). These polypeptides are either inducibly or constitutively expressed on various APCs and are membrane-bound ligands for CD28 and CTLA-4, respectively, on T-cells. CD28 (Aruffo and Seed Proc. Natl. Acad. Sci. 84, 8573-8577 (1987) and Gross et al. J. Immun. 144, 3201-3210 (1990)) is expressed on resting T-cells and mediates a positive costimulatory signal. CTLA-4 (Brunet et al. Nature 328, 267-270 (1987) and Dariavach et al. Eur. Jour. Immun. 18, 1901-1905 (1988)) expression is induced upon T-cell activation and negatively regulates the CD28 signal, due to its higher binding affinity for B7.1 and B7.2. Mice without the CTLA-4 gene exhibit dramatically high levels of T-cells, since the switch off mechanism for the proliferation signal is impaired in the absence of CTLA-4. This phenotype clearly demonstrates the major inhibitory effect that the CTLA-4 costimulatory protein has on T-cell proliferation. Mice lacking CD28 or B7.1 or B7.2 have a less severe phenotype, indicating that alternate pathways for T-cell costimulation may exist.

 There has been considerable interest in the CD28/CTLA-4 pathway as means for regulating T-cell activation and proliferation. A chimeric protein containing the extracellular portion of CTLA-4 fused to

- 3 -

human Fc has strong immunosuppressive effects and has been studied in a variety of clinical settings.

Antibodies to B7.1 and B7.2 proteins have also been evaluated for similar indications in the area of

5 immunosuppression. Anti-CTLA-4 antibodies have shown utility in promoting T-cell activation. In addition, B7.1 and B7.2 gene therapy has shown great promise in the area of cancer immunotherapy.

Thus far, CD28, CTLA-4, B7.1 and B7.2 are
10 involved in a single T-cell costimulatory pathway. Given the capability of modulating an immune response by regulating T-cell costimulation, it would be desirable to identify other members of the same or a separate T-cell costimulatory pathway which may have
15 advantageous properties in regulating host T-cell function and immune response.

Accordingly, it is an object of the invention to provide novel polypeptides for stimulation of T-cell activity and/or proliferation. It is a further object
20 of the invention to use the novel polypeptides for the prevention and treatment of T-cell mediated immune disorders.

Summary of the Invention

25 Surprisingly, two novel polypeptides of a T-cell costimulatory pathway have been identified. The polypeptides represent a ligand-receptor pair in a unique costimulatory pathway which appears to be distinct from the pathway consisting of previously
30 described proteins CD28, CTLA-4, B7.1, and B7.2. The polypeptides are referred to as CD28 related protein-1, or CRP1, and B7 related protein-1, or B7RP1.

The invention provides for nucleic acid molecules encoding CRP1 and B7RP1 polypeptides and
35 related polypeptides. An isolated nucleic acid

- 4 -

molecule of the invention comprises a nucleotide sequence selected from the group consisting of:

- a) the nucleotide sequence as set forth in Figure 1A (SEQ ID NO:1);
- 5 b) the nucleotide sequence encoding the polypeptide from residues 1-200 or from residues 21-200 as set forth in Figure 1A (SEQ ID NO:1);
- c) a nucleotide sequence encoding a polypeptide that is at least about 70 percent identical
- 10 to the polypeptide as set forth in Figure 1A (SEQ ID NO: 1);
- d) a naturally occurring allelic variant or alternate splice variant of any of (a), (b) or (c);
- e) a nucleotide sequence complementary to any
- 15 of (a), (b) or (c);
- f) a nucleotide sequence of (b), (c) or (d) encoding a polypeptide fragment of at least about 25, 50, 75, 100, or greater than 100 amino acid residues;
- g) a nucleotide sequence of (a), (b) or (c)
- 20 comprising a fragment of at least about 10, 15, 20, 25, 50, 75, 100, or greater than 100 nucleotides; and
- h) a nucleotide sequence which hybridizes under stringent conditions to any of (a)-(g).

25 Also provided by the invention is an isolated nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:

- a) the nucleotide sequence as set forth in Figure 2A (SEQ ID NO: 6) or Figure 3A (SEQ ID NO:11) or
- 30 Figure 12A (SEQ ID NO: 16);
- b) the nucleotide sequence encoding the polypeptide as set forth in Figure 2A (SEQ ID NO: 6) from residues 1-322 or from residues 47-322 or as set forth in Figure 3A (SEQ ID NO:11) from residues 1-288
- 35 or from residues 19-288, 20-288, 21-288, 22-288, 24-288, or 28-288; or as set forth in Figure 12A from

- 5 -

residues 1-302 or from residues 19-302, 20-302, 21-302, 22-302, 24-302 or 28-302;

5 c) a nucleotide sequence encoding a polypeptide that is at least about 70 percent identical to the polypeptide as set forth in Figure 2A (SEQ ID NO: 6) or Figure 3A (SEQ ID NO: 11) or Figure 12A (SEQ ID NO: 16);

d) a naturally occurring allelic variant or alternate splice variant of any of (a), (b) or (c);

10 e) a nucleotide sequence complementary to any of (a), (b) or (c);

f) a nucleotide sequence of (b), (c) or (d) encoding a polypeptide fragment of at least about 25, 50, 75, 100, or greater than 100 amino acid residues;

15 g) a nucleotide sequence of (a), (b) or (c) comprising a fragment of at least about 10, 15, 20, 25, 50, 75, 100, or greater than 100 nucleotides; and

h) a nucleotide sequence which hybridizes under stringent conditions to any of (a)-(g).

20

The subject matter of the invention also relates to CRP1 and B7RP1 polypeptides and related polypeptides. The invention provides for an isolated polypeptide comprising the amino acid sequence selected from the group consisting of:

25 a) the amino acid sequence as set forth in Figure 1A (SEQ ID NO:2);

30 b) the mature amino acid sequence as set forth in Figure 1A (SEQ ID NO:2) comprising a mature amino terminus at residue 21;

c) a fragment of the amino acid sequence set forth in Figure 1A (SEQ ID NO: 2) comprising at least about 25, 50, 75, 100, or greater than 100 amino acid residues;

35 d) an ortholog of (a), (b) or (c); and

- 6 -

e) an allelic variant or alternative splice variant of (a), (b) or (d).

Also in accordance with the invention is an
5 isolated polypeptide comprising the amino acid sequence selected from the group consisting of:

a) the amino acid sequence as set forth in Figure 2A (SEQ ID NO: 7) or Figure 3A (SEQ ID NO:12) or Figure 12A (SEQ ID NO: 17);

10 b) the mature amino acid sequence as set forth in Figure 2A (SEQ ID NO: 7) comprising a mature amino terminus at residues 47 or Figure 3A (SEQ ID NO:12) comprising a mature amino terminus at any of residues 19, 20, 21, 22, 24 or 28 or Figure 12A (SEQ ID
15 NO: 17) comprising a mature amino terminus at any of residues 19, 20, 21, 22, 24, or 28;

c) a fragment of the amino acid sequence set forth in Figure 2A (SEQ ID NO: 7) or Figure 3A (SEQ ID NO: 12) or Figure 12A (SEQ ID NO: 17) comprising at
20 least about 25, 50, 75, 100, or greater than 100 amino acid residues;

d) an ortholog of (a), (b) or (c); and

e) an allelic variant or alternative splice variant of (a), (b), (c) or (d).

25

Also encompassed by the invention are expression vectors and host cells for production of the polypeptides, antibodies which bind to CRP1 and B7RP1 polypeptides and to related polypeptides, and assays
30 for detecting binding of B7RP1 and B7RP1-related polypeptides to CRP1 and CRP1-related polypeptides. Pharmaceutical compositions comprising CRP1 or CRP1-related polypeptides and B7RP1 or B7RP1-related polypeptides are also encompassed by the invention.
35 Methods for identifying compounds that interact with CRP1 or B7RP1 are also provided as are assays for

- 7 -

determining whether such compounds are agonists or antagonists of CRP1 and B7RP1 activity.

CRP1 and B7RP1 polypeptides are involved in T-cell costimulation and proliferation. CRP1 and B7RP1
5 polypeptides, selective binding agents thereof, and agonists and antagonists thereof, may be useful for the diagnosis, prevention and treatment of diseases related to the control of T-cell responses.

CRP1 and B7RP1 polypeptides, selective
10 binding agents thereof, and agonists and antagonists thereof, may be useful for the diagnosis, prevention and treatment of immune disorders, either for stimulating insufficient immune response or reducing or inhibiting an exaggerated or inappropriate immune
15 response. The immune disorder may be mediated directly or indirectly by T cells.

The invention provides for treating, preventing, or ameliorating a T-cell mediated disorder comprising administering to an animal a CRP1 or B7RP1
20 polypeptide. The invention also provides for a method of diagnosing a T-cell mediated disorder or a susceptibility to a T-cell mediated disorder in an animal comprising determining the presence or amount of expression of a CRP1 or B7RP1 polypeptide; and
25 diagnosing a T-cell mediated disorder or a susceptibility to a T-cell mediated disorder based on the presence or amount of expression of the polypeptide. Typically, a T-cell mediated disorder is an immune disorder which may be mediated directly or
30 indirectly by T cells. The animal is preferably a mammal and more preferably a human. The invention also provides for a method of identifying a test molecule which binds to a CRP1 or B7RP1 polypeptide comprising contacting the polypeptide with a test compound and
35 determining the extend of binding of the polypeptide to the test compound. The method may be used to identify

- 8 -

agonists and antagonists of CRP1 and/or B7RP1 polypeptide.

Antagonists of CRP1 and/or B7RP1 polypeptides may be used as immunosuppressive agents for many indications, including autoimmune disorders (such as rheumatoid arthritis, psoriasis, multiple sclerosis, diabetes, and systemic lupus erythematosus), toxic shock syndrome, bone marrow and organ transplantation, inflammatory bowel disease, allosensitization due to blood transfusions, and the treatment of graft vs. host disease. In addition, antagonists may be used as inhibitory agents for T-cell dependent B-cell mediated indications including asthma and allergy, and antibody mediated autoimmunity. Agonists of the CRP1 and/or B7RP1 polypeptides may be useful in, but not restricted to, T-cell activation for tumor surveillance and removal.

An antagonist of the invention includes an antibody, or fragment thereof, which is reactive with or binds to B7RP1 or to an extracellular domain of B7RP1 wherein the antibody reduces or eliminates the binding to B7RP1 to CRP1. In one embodiment, the antibody binds selectively to human B7RP1 or to an extracellular domain thereof. The antibody or fragment thereof which is an antagonist inhibits partially or completely the immune costimulatory activity of B7RP1. In a preferred embodiment, the antibody is a monoclonal antibody and may be murine, human, chimeric or humanized.

The invention further provides for a method of regulating the interaction of B7RP1 with CRP1 comprising administering to an animal a selective binding agent of CRP1 or a selective binding agent of B7RP1 or both. In one embodiment, the selective binding agent is an antibody which binds to B7RP1 and reduces or eliminates the binding to B7RP1 to CRP1.

- 9 -

The invention also provides a method of regulating immune costimulation mediated by B7RP1 comprising administering to an animal a selective binding agent of B7RP1. The selective binding agent is preferably an antibody which binds to B7RP1 and partially or completely inhibits immune costimulation mediated by B7RP1.

The invention also provides for a method of regulating T-cell activation or proliferation in an animal comprising administering to the animal a nucleic acid molecule encoding a CRP1 or B7RP1 polypeptide. For example, a nucleic acid molecule encoding a B7RP1 polypeptide may be used in gene therapy to enhance T-cell activation in response to various tumors.

Also encompassed by the invention is a transgenic non-human mammal comprising a nucleic acid molecule encoding a CRP1 or B7RP1 polypeptide. The CRP1 or B7RP1 nucleic acids are introduced into the mammal in a manner that allows expression and increased circulation levels of CRP1 or B7RP1 polypeptides. The transgenic non-human mammal is preferably a rodent, and more preferably a mouse or a rat.

Description of Figures

Figure 1. A) DNA and amino acid sequence murine CRP1 (mCRP1). Predicted signal sequence of CRP1 is underlined at the amino-terminus and the experimentally determined pro-peptide cleavage site is indicated by an asterisk. Predicted transmembrane sequence is underlined toward the carboxy-terminus. B) Amino acid alignment of murine CRP1 protein sequence (mCRP1) with murine CD28 (mCD28).

Figure 2. A) DNA and amino acid sequence of murine B7RP1 (mB7RP1). Predicted signal sequence of B7RP1 is underlined at the amino-terminus and the

- 10 -

experimentally determined pro-peptide cleavage site is indicated by an asterisk. Predicted transmembrane sequence is underlined toward the carboxy-terminus. B) Amino acid alignment of B7RP1 protein sequence (mB7RP1) with murine CD80 (mCD80).

Figure 3. A) Structure and sequence of the protein coding region of the putative human B7RP1 (hB7RP1). Predicted signal sequence of hB7RP1 is underlined at the amino-terminus. Predicted signal peptide cleavage sites are marked by asterisks. Predicted transmembrane sequence is underlined toward the carboxy-terminus. B) Amino acid alignment of the putative mature hB7RP1 protein with the mature murine B7RP1 (mB7RP1) protein.

Figure 4. A) Expression of soluble CRP1-Fc fusion protein from 293T cells transfected with the pCDNA3/CRP1-Fc. Normalized volumes of cell lysate or conditioned medium were loaded and separated on a 10% PAGE gel as indicated. Western analysis of cell lysate and cell media supernatant for expression of cell-associated (cell lysate) and secreted (media) Fc fusion proteins. Primary antibody was Goat-anti human Fc antibody (Pierce Chemical Company, Rockford, IL.). B) Expression of soluble B7RP1-Fc fusion protein from 293T cells transfected with the pCDNA3/ B7RP1-Fc. 20 µl of normalized cell lysate or media supernatant were loaded and separated on a 10% PAGE gel. Western analysis was conducted as in (A).

Figure 5. Interaction of CRP1-Fc and B7RP1-Fc fusion proteins with membrane-bound proteins expressed in COS-7 cells. COS-7 cells transiently transfected with pCDNA3/CRP1, pCDNA3/B7RP1, or pCDNA3 vector alone. CHO D-cells were transfected with

- 11 -

psDR α /hCD28 and stably expressed human CD28 (hCD28). Cells expressing membrane-bound CRP1, B7RP1, or hCD28, are represented in rows as indicated at the left side of the panel. Fc fusion proteins were incubated with
5 the plate-bound cells in columns as indicated at the top of the panel. After incubation, cells were washed, and bound Fc fusion proteins were detected using an anti-human Fc antibody and ACAS (Adherent Cell Analysis and Sorting; ACAS Ultima, Meridian Instruments, Inc.,
10 Okemos, MI) analysis.

Figure 6. FACS (Fluorescence-Activated Cell Sorter) analysis of expression of the receptor for B7RP1 (putatively, CRP1) on activated CD4+ and CD8+
15 T-cells. Mouse splenocytes were activated with PMA and ionomycin for 12 hours. B7RP1-Fc fusion protein, control Fc protein (Mock-Fc), or PBS (no stain), were incubated with the cells, washed, and subsequently incubated with goat-anti-human Fc-FITC conjugated
20 antibody (GaHuFc-FITC) as indicated at the bottom of each panel. Cell marker antibodies (for T-cell markers CD4 and CD8) PE conjugated, or isotype control antibody (rat isotype) PE conjugated, or PBS (no stain), were added as indicated at the left side of each individual
25 panel.

Figure 7. FACS (Fluorescence-Activated Cell Sorter) analysis of the expression of B7RP1 on B-cells. Fluorocytometric analysis of the expression of the
30 ligand for CRP1 (presumably, B7RP1) on mouse splenocytes. CRP1-Fc fusion protein, control Fc protein (Mock-Fc), or PBS (no stain), were incubated with the cells, washed, and subsequently incubated with goat-anti-human Fc-FITC conjugated antibody
35 (GaHuFc-FITC) as indicated at the bottom of each panel.

- 12 -

PE conjugated cell marker antibody to CD45R (CD45R is a B-cell marker) or isotype control antibody (rat isotype), or PBS (no stain), were added as indicated at the left side of each individual panel.

5

Figure 8. FACS analysis of the expression of mCRP1 ligand on peritoneal macrophages. Peritoneal cells were first distinguished in subsets on the ground of their light scattering properties (panel A).

10 Macrophages were identified in region 5 (R5) because of their ability to strongly scatter light forward (FSC) and sideways (SSC) and because of their positive staining for the F4/80 antigen, a marker for macrophages (panel B). Macrophages in region 6 (R6)
15 were singled out on the basis of their less intense staining for the F4/80 antigen and found to be stained by the CRP1-Fc fusion protein (presumably because of their expression of B7RP1).

20

Figure 9. Inhibition of T-cell proliferation using a B7RP1-Fc fusion protein. T-cells from mouse splenocytes were activated by increasing concentrations of Concanavalin A (Con A) as indicated at the bottom of the graph. mCRP1-Fc, mB7RP1, and mB7.2-Fc fusion

25

proteins were added to enriched T-cells from splenocytes in the absence (no adds) or presence of Con A. 200,000 cells were used in the T-cell proliferation assays in a 96-well plate. Cells were incubated with media (no adds) or Fc fusion proteins as indicated in
30 the graph legend. After 42 hr, cells were pulsed with H-thymidine for 6 hr, then harvested and incorporated radioactivity determined. Average CPM and standard deviation from triplicate samples are represented.

35

Figure 10. A) Normal mesenteric lymph node from control Mouse #10 showing the cortex, paracortex

- 13 -

and medulla of the node. Hematoxylin-eosin (H&E) stain, 40x magnification. B) Markedly enlarged mesenteric lymph node from WX11 Mouse #40 with prominent follicular hyperplasia (FH), expansion of paracortex and medullary cord hyperplasia (MH). H&E, 40x. C) Close-up of the medullary cords (MC) and sinuses (MS) from the mesenteric lymph node of control Mouse #10. Note the small medullary cords composed of mostly small lymphocytes adjacent to medullary sinuses with fleshy macrophages. H&E, 400x. D. Close-up of the medullary cords (MC) and sinuses (MS) from the mesenteric lymph node of WX11 Mouse #40. Note the markedly thickened medullary cords composed of large numbers of plasma cells with occasional Russell body cells (arrow). H&E, 400x. E) Normal spleen from control Mouse #10 showing red pulp and white pulp areas with periarteriolar lymphoid sheaths (PALS), 100x. Inset: close-up of the marginal zone surrounding the white pulp with small lymphocytes, macrophages and occasional plasma cells, 400x. F) Spleen from WX11 Mouse #6 with enlarged white pulp areas, including PALS and follicles (arrow), 100x. Inset: close-up of the marginal zone with numerous plasma cells and occasional Russell bodies, 400x. G) Ileum with Peyer's patch from control Mouse #25 with the interfollicular zone (arrow) flanked by two secondary follicles, 40x. H) Ileum with Peyer's patch from WX11 Mouse #32 with markedly enlarged follicles with prominent germinal centers and interfollicular tissue (arrow), 40x.

30

Figure 11. A) Normal mesenteric lymph node from control Mouse #5 showing the cortex, paracortex and medulla of the node. Hematoxylin-eosin (H&E) stain, 40x magnification. B) Markedly enlarged mesenteric lymph node from WX11 Mouse #33 with prominent follicular hyperplasia (top: rows of secondary

35

- 14 -

follicles in the outer cortex), expansion of the paracortex (center) and medullary cord hyperplasia (bottom). H&E, 40x. C) Immunohistochemical staining of the mesenteric lymph node from control Mouse #10 with anti-B220 antibody (B cell marker). Note the intensely (brown) staining cortical area and thin medullary cords. Immunostaining performed using the avidin-biotin complex (ABC) immunoperoxidase method (DAB chromogen, hematoxylin counterstain), 40x. D) Immunohistochemical staining of the mesenteric lymph node from WX11 Mouse #33 with anti-B220 antibody. Note the intensely staining cortical follicles and medullary cords (although the mature plasma cells in the cords are negative for B220), 40x. E) Immunohistochemical staining of the lymph node from control Mouse #10 with anti-CD3 antibody (T-cell marker). Note the immunostaining of the paracortical zone of the node, 40x. F) Immunohistochemical staining of the lymph node from WX11 Mouse #33 with anti-CD3 antibody. Note the enlarged, intensely staining paracortical areas of the node, 40x.

Figure 12. A) Structure and sequence of the protein coding region of human B7RP1 (hB7RP1). Predicted signal sequence of hB7RP1 is underlined at the amino-terminus. Predicted signal peptide cleavage sites are marked by asterisks. Predicted transmembrane sequence is underlined toward the carboxy-terminus. B) Amino acid alignment of the putative mature hB7RP1 protein with the mature murine B7RP1 (mB7RP1) protein.

Figure 13. A) Structure and sequence of the protein coding region of human CRP1 (hCRP1). Predicted signal sequence of hCRP1 is underlined at the amino-terminus. Predicted signal peptide cleavage sites are

- 15 -

marked by asterisks. Predicted transmembrane sequence is underlined toward the carboxy-terminus. B) Amino acid alignment of the hCRP1 protein with the murine CRP1 (mB7RP1) protein.

5

Figure 14. CRP-1 is on resting memory T-cells. Resting splenocytes from 6-7 month old mice were double-stained using B7RP-1-Fc labeled by an FITC-conjugated anti-human Fc antibody and a PE-conjugated antibody to either CD44(Fig 14A), CD45RB(Fig 14B), or CD69(Fig 14C).

Figure 15. T-cell co-stimulation by B7RP-1-Fc fusion protein. A) T-cell proliferation induced by different quantities of B7RP-1-Fc (closed squares), B7.2-Fc (closed circles), or OPG-Fc fusion protein control (open squares) in conjunction with anti-CD3 antibody. Fusion proteins were used at various concentrations to coat 96 well plates pre-coated with anti-human Fc FAb₂ (12.5 µg/ml) and anti-CD3 antibody (0.9 µg/ml). B7RP-1-Fc and B7.2-Fc co-stimulate T-cells in a dose-dependent fashion up to 0.3 µg/ml, at which the maximal effect is achieved. B) T-cell proliferation induced by B7RP-1-Fc (closed squares), B7.2-Fc (closed circles), non-fused Fc (open squares), or no Fc (open circles) in conjunction an anti-CD3 antibody (0.85 µg/ml) and in the presence of various concentrations of a rabbit anti-B7RP-1-Fc polyclonal antibody. Fc fusion proteins were used at a concentration of 0.3 µg/ml and were bound to the plates as above. The anti-B7RP-1-Fc antibody was raised to purified B7RP-1-Fc by subcutaneous injections of antigen emulsified in adjuvant, and then was affinity purified. The antibodies were incubated for 30 min with the Fc fusion proteins before the addition of the cells. The anti-B7RP-1-Fc antibody specifically

- 16 -

inhibits the T-cell proliferation induced by B7RP-1-Fc in a dose-dependent fashion.

Figure 16. Effect of CRP-1-Fc and B7RP-1-Fc proteins on the incidence (A) and severity (B) of collagen induced arthritis in mice. Collagen induced arthritis susceptible B10.RIII mice were immunized at the base of the tail with 10 µg porcine collagen type II in CFA. Mice received 100 µg of fusion protein twice per week. Fc fusion proteins and control PBS treatment are indicated in the figure legend.

Figure 17. Proximal Colon in B7RP-1-Fc Transgenic Mice. (A) Normal proximal colon from control Mouse#53F (female) showing the gut wall with mucosa, submucosa, muscularis and serosa. Hematoxylin-eosin (H&E) stain, 40x magnification. (B) Diffusely thickened proximal colon from B7RP-1-Fc transgenic Mouse#111F with prominent glandular hypertrophy, fissuring ulceration and transmural inflammation. H&E, 40x. (C) Lower power view of proximal colon (as in panel B) from B7RP-1-Fc transgenic Mouse#111F with multifocal fissuring ulceration and transmural inflammation. H&E, 20x. (D) Close-up of the fissuring ulcer and hypertrophic colonic glands from B7RP-1-Fc transgenic Mouse#111F (shown in panels B and C above). Note the lumen with mucopurulent exudate. H&E, 100x. (E) Close-up of granulomatous inflammation in the submucosa of B7RP-1-Fc transgenic Mouse#112F with a multinucleated giant cell surrounded by macrophages, lymphocytes and fewer neutrophils. H&E, 400x. (F) Close-up of granulomatous inflammation in the mucosa of B7RP-1-Fc transgenic mouse #112F with epithelioid macrophages mixed with lymphocytes, plasma cells and

- 17 -

fewer neutrophils subjacent to mucosal glands. H&E, 400x.

Figure 18. Distal Colon in B7RP-1-Fc Transgenic Mice. (A) Normal distal colon from control Mouse#53F (female) showing the layers of the gut wall with mucosa, submucosa, muscularis and serosa. Hematoxylin-eosin (H&E) stain, 40x magnification. (B) Diffusely thickened distal colon from B7RP-1-Fc transgenic Mouse#111F (female) with prominent glandular hypertrophy and hyperplasia and scattered crypt abscesses. H&E, 40x. (C) Diffusely thickened distal colon from B7RP-1-Fc transgenic Mouse#55M (male) with prominent glandular hypertrophy and hyperplasia. H&E, 40x. (D) Diffusely thickened distal colon from B7RP-1-Fc transgenic Mouse#112F (female) with hypertrophic colonic glands, focal lymphoid aggregates and many crypt abscesses. H&E, 40x. (E) Immunohistochemical staining of the distal colon from B7RP-1-Fc transgenic Mouse#112F with anti-CD3 antibody (T cell marker). Note the immunostaining of the superficial mucosa and colonic lymphoid patch. H&E, 40x. (F) Close-up of the colonic mucosa of B7RP-1-Fc transgenic Mouse#112F with a crypt abscess (arrow) and lymphoid aggregate composed of B220+ B cells (inset). H&E, 100x.

Figure 19. Small Intestine in B7RP-1-Fc Transgenic Mice. (A) Normal duodenum from control Mouse#53F (female) showing the lumen, villi and crypts of the mucosa and underlying submucosa, muscularis and serosa. Hematoxylin-eosin (H&E) stain, 40x magnification. (B) Diffusely thickened duodenum from B7RP-1-Fc transgenic Mouse#51F (female) with prominent crypt hypertrophy and hyperplasia and mild lymphoplasmacytic infiltrate in the lamina propria. H&E, 40x. (C) Normal jejunum from control mouse #53F

- 18 -

(female) showing the normal length of the villi and crypts in the jejunal mucosa. H&E, 40x magnification. (D) Markedly thickened jejunal mucosa from B7RP-1-Fc transgenic Mouse#51F (female) with locally extensive crypt hypertrophy and hyperplasia. H&E, 40x. (E) Normal ileum from control Mouse#53F (female) showing the normal length of the villi and crypts in the ileal mucosa. H&E, 40x magnification. (F) Mild atrophy of ileal mucosa from B7RP-1-Fc transgenic Mouse#231M (male) with focal loss and blunting of villi. H&E, 40x.

Figure 20. The B7RP-1-Fc fusion protein inhibits tumor growth in mice. Meth A sarcoma cells were implanted intradermally in the abdomen of Balb/C mice. On days 7, 10, 14, and 17, after implantation, the mice were treated with vehicle (dark diamonds) or murine B7RP-1-Fc (grey triangles, Example 7). Tumor volume was measured, as described in Example 20, on the indicated days after implantation. The tumor growth was monitored up to day 28. Each group had eight mice.

Figure 21. T-cell co-stimulation by human B7RP-1-Fc. Anti-CD3 and human B7RP-1 Fc were used to coat 96 well plates, and 1×10^5 T-cells/well (>98% CD3+) were cultured and harvested as described in Example 21. A) Co-stimulation induced by anti-CD3 only (closed circles), 0.5 μ g/ml B7RP-1 Fc (closed triangles), 0.5 μ g/ml OPG-Fc (open circles), and 5 μ g/ml anti-CD28 (open triangles) at different concentrations of anti-CD3 primary stimulation. Data show that B7RP-1-Fc co-stimulated anti-CD3 primed T-cells to similar levels as co-stimulation using anti-CD28 antibodies. Data shown are mean [3 H]TdR incorporated \pm SD in triplicate wells from one representative experiment of several experiments

- 19 -

generated with T-cells isolated from three normal donors. B) Dose-dependent inhibition of B7RP-1-Fc co-stimulation by CRP-1-Fc. T-cells were cultured in wells coated with both anti-CD3 at 0.3 µg/ml and 0.5 µg/ml B7RP-1-Fc. Serially diluted concentrations of CRP-1-Fc (closed circles) or OPG-Fc (open circles) were preincubated with the B7RP-1-Fc for 30 min prior to the addition of T-cells. Data show that CRP-1-Fc inhibits B7RP-1 induced co-stimulation in a dose-dependent manner. Percent inhibition is plotted against CRP-1-Fc or OPG-Fc protein concentration. Data shown are mean [³H]TdR incorporated +/- SD of three experiments done in triplicate wells and are representative of experiments generated with two normal donors. C) Co-stimulation by CHO human B7RP-1 cells. T-cells were purified from peripheral blood and were cultured with various concentrations of anti-CD3 in the presence of anti-CD3 alone (closed circles), 1 x 10⁴ CHO vector control cells (open circles) or 1 x 10⁴ CHO B7RP-1 cells (closed triangles), as described in Example 22. The data show that membrane-bound B7RP-1 co-stimulated T-cell growth to a level similar to that observed using B7RP-1-Fc fusion proteins. Data shown are the mean +/- SD of triplicate cultures and are representative of results generated with two normal donors. D) Cytokine production. T-cells were cultured as described in (Figure 21A) and supernatants were collected at 48 (black bars) and 72 (gray bar) hrs. Data show that the amount of IL-2 produced by B7RP-1-Fc co-stimulated cells (top graph) was similar to that produced by cells stimulated by anti-CD3 and control Fc, but significantly less than that produced by anti-CD28 co-stimulated cells. Data also show that B7RP-1-Fc co-stimulation enhanced IL-10 (middle graph) and IFN-gamma (bottom graph) production.

- 20 -

Detailed Description of the Invention

The invention provides for novel polypeptides referred to herein as CRP1 and B7RP1, which comprise a receptor-ligand pair that is involved in T-cell activation. cDNAs encoding the polypeptides were identified from a library prepared from mouse intestinal intraepithelial cells and screened on the basis of homology to the CD28 and CTLA-4 polypeptides (for CRP1) or B7.1 and B7.2 polypeptides (for B7RP1).

CD28 related protein-1, or CRP1, is predicted to be a type I transmembrane protein with a signal sequence and extracellular domain at the amino-terminus, a transmembrane domain, and a carboxy terminal intracellular domain (Figure 1). The full-length CRP1 protein is 180 amino acids in its mature form. The predicted leader sequence spans about amino acid residues 1-20 (relative to the initiating methionine) and the extracellular domain of the mature protein encompasses about residues 21-145 (Example 1). The predicted transmembrane domain spans about residues 146-163 and the intracellular domain encompasses about residues 164-200. The amino terminal extracellular domain is similar to an Ig loop with conserved putative intra- and inter-molecular bonding cysteines. Furthermore, a "MYPPPY" motif, which is previously known to be important for B7.1 and B7.2 binding to CD28 and CTLA-4, is also partially conserved.

CD28 and CTLA-4 are weakly homologous as exemplified by the 26% amino acid identity between murine CD28 and CTLA-4. There is 19% amino acid identity of CRP1 with murine CD28 and 14% identity of CRP1 with murine CTLA-4. However, critical cysteine residues are conserved between murine CD28, CTLA-4 and CRP1 at residues 42, 63, 83, 109, and 137 (relative to

- 21 -

the initiating methionine in the CRP1 protein, See Figure 1A). The approximate mature protein lengths and locations of the transmembrane region relative to the carboxy terminus are also similar in CRP1, CD28, and CTLA-4.

Human CRP1 is a transmembrane protein having the nucleotide and amino acid sequence as shown in Figure 13A. The predicted leader sequence spans about residues 1-19 or about residues 1-20. The predicted mature amino terminus is at residues 20 or 21. Preferably, the mature amino terminus is at position 21. The extracellular domain spans from any of the predicted mature amino termini to about amino acid residue 140, the transmembrane domain spans about residues 141-161 and the intracellular domain spans about residues 162-199. Human CRP1 protein has 69% identity to the murine protein and the corresponding nucleotide sequences are 77% identical. The sequence of human CRP1 was reported in Hutloff et al. Nature 397, 263-266 (1999).

B7 related protein-1, or B7RP1, is predicted to be a type I transmembrane protein with a signal sequence and extracellular domain at the amino-terminus, a transmembrane domain, and a carboxy terminal intracellular domain (Figure 2A). The full-length B7RP1 protein is 276 amino acids in its mature form. The predicted leader sequence spans about amino acid residues 1-46 (relative to the initiating methionine) and the extracellular domain of the mature protein encompasses residues 47-279 (Example 3). The predicted transmembrane domain spans residues 280-298 and the intracellular domain encompasses residues 299-322. Similar to B7.1 and B7.2, the extracellular domain of B7RP1 comprises two Ig loops.

B7.1 and B7.2 are weakly homologous as exemplified by the 24% amino acid identity between

- 22 -

murine B7.1 and B7.2. There is 20% amino acid identity of B7RP1 with murine B7.1 and 19% identity of B7RP1 with murine B7.2. However, critical cysteine residues are conserved between murine B7.1, B7.2 and B7RP1 at residues 62, 138, 185, and 242 (relative to the initiating methionine in the B7RP1 protein, Figure 2A). The approximate mature protein length and location of the transmembrane region relative to the carboxy terminus are also similar in mB7RP1, B7.1, and B7.2.

Human B7RP1 is also a transmembrane protein with conserved cysteine residue in the extracellular domain which are necessary for Ig loop structures. The predicted leader sequence encompasses about residues 1-18, 1-19, 1-20, 1-21, 1-23 or 1-27 as shown in Figure 3A. The predicted mature amino terminus may be at any of the residues 19, 20, 21, 22, 24 or 28. Preferably, the amino terminus is at position 19. The extracellular domain spans from any of the mature amino termini to about amino acid residue 259. The predicted transmembrane domain spans about residues 259-274. The intracellular domain encompasses residues 275-302. The full-length human B7RP1 nucleotide and amino acid sequence is shown in Figure 12A. Human B7RP1 is about 43% identical to the murine protein.

CRP1 and B7RP1 bind each other, but CRP1 does not detectably bind to the B7RP1 related protein B7.2; and B7RP1 does not exhibit detectable binding to CRP1-related CD28 or CTLA-4 (Example 8). B7RP1 was shown to regulate T-cell proliferation, presumably through the interaction of B7RP1 with CRP1 receptors (Example 11). Thus, CRP1 and B7RP1 represent a novel pathway for regulating T-cell proliferation and activation.

The interaction of B7RP1 with CRP1 can be regulated in such a manner that immune costimulation and T-cell proliferation and activation can be

- 23 -

increased or decreased. By way of example, anti-B7RP1 monoclonal and polyclonal antibodies raised against murine B7RP1 blocked the B7RP1/CRP1 interaction and also blocked T-cell proliferation induced by a B7RP1-Fc fusion protein (see Example 17). A human CRP-1-Fc fusion protein blocked human T-cell proliferation induced by human B7RP-1-Fc (Example 21). In addition, addition of a CRP1-Fc fusion protein delayed the onset of arthritic symptoms in a mouse model of rheumatoid arthritis (see Example 18). B7RP-1/CRP-1 co-stimulation can also be increased by addition of B7RP-1-Fc fusion protein or other activators of this pathway (Example 20).

15

Nucleic Acid Molecules

The term "isolated nucleic acid molecule" refers to a nucleic acid molecule that is free from at least one contaminating nucleic acid molecule with which it is naturally associated, and preferably substantially free from any other contaminating mammalian nucleic acid molecules.

The term "allelic variant" refers to one of several possible naturally occurring alternate forms of a gene occupying a given locus on a chromosome of an organism.

The term "splice variant" refers to a nucleic acid molecule, usually RNA, which is generated by alternative processing of intron sequences in an RNA transcript.

The term "high stringency conditions" refers to those conditions which: (1) employ low ionic strength and high temperature for washing, for example, 0.1 X SSC (0.015 M NaCl/0.0015 M sodium citrate) 0.1% NaDodSO₄ (SDS) at 50°C, or (2) employ during

- 24 -

hybridization a denaturing agent such as formamide, for example, 50% (vol/vol) formamide with 0.1% bovine serum albumin/0.1%. Ficoll/0.1% polyvinylpyrrolidone/50 mM sodium phosphate buffer at pH 6.5 with 5 X SSC (750 mM NaCl, 75 mM sodium citrate) at 42°C. Another example of high stringency conditions is 50% formamide, 5 x SSC, 50 mM sodium phosphate (pH 6.8), 0.1% sodium pyrophosphate, 5 x Denhardt's solution, sonicated salmon sperm DNA (50 µg/ml), 0.1% SDS, and 10% dextran sulfate at 42°C, with washes at 42°C in 0.2 x SSC and 0.1% SDS.

The term "moderate stringency conditions" refers to those conditions which include the use of a washing solution and hybridization conditions (e.g., temperature and ionic strength) less stringent than described above. An example of moderately stringent conditions are conditions such as overnight incubation at 37°C in a solution comprising: 20% formamide, 5 x SSC, 50 mM sodium phosphate (pH 7.6), 5 x Denhardt's solution, 10% dextran sulfate, and 20 µl/ml denatured sheared salmon sperm DNA, followed by washing the filters in 1 x SSC at about 37-50°C. The skilled artisan will recognize how to adjust the temperature, ionic strength and other parameters as necessary to accommodate factors such as probe length and the like.

Recombinant DNA technology methods are set forth in Sambrook et al. (*Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY (1989)) and/or Ausubel et al., eds., (*Current Protocols in Molecular Biology*, Green Publishers Inc. and Wiley and Sons, NY (1994)) which are hereby incorporated by reference in their entirety.

- 25 -

The invention provides for isolated nucleic acid molecules encoding CRP1 and B7RP1 polypeptides. Also provided for are nucleic acid molecules which are fragments, allelic variants, splice variants, or are
5 complementary in sequence to molecules encoding CRP1 and B7RP1 polypeptides. Nucleic acid molecules which are at least about 70% identical to molecules encoding CRP1 or B7RP1 or which hybridize to molecules encoding CRP1 or B7RP1 under moderate or high stringency
10 conditions are also encompassed. The nucleic acid molecules may be cDNA, genomic DNA, RNA or a partially or totally synthetic nucleic acid molecule. In preferred embodiments, nucleic acid molecules of the invention are at least about 75%, 80%, 85%, 90% or 95%
15 identical to nucleic acid molecules encoding CRP1 or B7RP1.

A gene or cDNA encoding a CRP1 or B7RP1 polypeptide or fragment thereof may be obtained, for example, by hybridization screening or PCR
20 amplification of a genomic or cDNA library. Probes or primers useful for screening the library can be generated based on sequence information for other known genes or gene fragments from the same or a related family of genes, for example, conserved motifs found in
25 CRP1 or B7RP1 related polypeptides such as a conserved array of cysteine residues. In addition, where a gene encoding a CRP1 or B7RP1 polypeptide has been identified from one species, all or a portion of that gene may be used as a probe to identify homologous
30 genes from other species. The probes or primers may be used to screen cDNA libraries from various tissue sources believed to express the CRP1 or B7RP1 gene.

Where oligonucleotide probes are used to screen cDNA or genomic libraries, one of the following
35 two high stringency solutions may be used. The first of these is 6 X SSC with 0.05% sodium pyrophosphate at

- 26 -

35°C-62°C, with the temperature depending on the length of the oligonucleotide probe. For example, 14 base pair probes are washed at 35-40°C, 17 base pair probes are washed at 45-50°C, 20 base pair probes are washed at 52-57°C, and 23 base pair probes are washed at 57-63°C. The temperature can be increased 2-3°C where the background non-specific binding appears high. A second high stringency solution utilizes tetramethylammonium chloride (TMAC) for washing oligonucleotide probes.

One stringent washing solution is 3 M TMAC, 50 mM Tris-HCl, pH 8.0, and 0.2% SDS. The washing temperature using this solution is a function of the length of the probe. For example, a 17 base pair probe is washed at about 45-50°C.

Another means to prepare a gene encoding a CRP1 or B7RP1 polypeptide or fragment thereof is to employ chemical synthesis using methods well known to the skilled artisan such as those described by Engels et al. (*Agnew. Chem. Intl. Ed.*, 28:716-734 (1989)). These methods include, *inter alia*, the phosphotriester, phosphoramidite, and H-phosphonate methods for nucleic acid synthesis. A preferred method for such chemical synthesis is polymer-supported synthesis using standard phosphoramidite chemistry. Typically, DNA encoding a CRP1 or B7RP1 polypeptide will be several hundred nucleotides in length. Nucleic acids larger than about 100 nucleotides can be synthesized as several fragments using these methods. The fragments can then be ligated together to form a full length CRP1 or B7RP1 polypeptide. Usually, the DNA fragment encoding the amino terminus of the polypeptide will have an ATG, which encodes a methionine residue. This methionine may or may not be present on the mature form of a CRP1 or B7RP1 polypeptide, depending on whether the polypeptide produced in the host cell is designed to be secreted from that cell.

- 27 -

CRP1 or B7RP1 nucleic acid molecules, fragments, and/or derivatives that do not themselves encode polypeptides that are biologically active may nonetheless be useful as hybridization probes in
5 diagnostic assays to test, either qualitatively or quantitatively, for the presence of CRP1 or B7RP1 DNA or corresponding RNA in mammalian tissue or bodily fluid samples.

In some cases, it may be desirable to prepare
10 nucleic acid and/or amino acid variants of naturally occurring CRP1 or B7RP1 polypeptides. Nucleic acid variants may be produced using site directed mutagenesis, PCR amplification, or other appropriate methods, where the primer(s) have the desired point
15 mutations (see Sambrook et al., *supra*, and Ausubel et al., *supra*, for descriptions of mutagenesis techniques). Chemical synthesis using methods described by Engels et al., *supra*, may also be used to prepare such variants. Other methods known to the
20 skilled artisan may be used as well.

Preferred nucleic acid variants include those containing codons which have been altered for optimal expression of CRP1 and B7RP1 polypeptides in a given host cell. Particular codon alterations will depend
25 upon the selection of protein and host cell. Such "codon optimization" can in one instance, be carried out by selecting codons which are preferentially used in highly expressed genes in a given host cell. Computer algorithms which incorporate codon frequency
30 tables such as "Ecohigh. Cod" for codon preference of highly expressed bacterial genes may be used and are provided by the University of Wisconsin Package Version 9.0, Genetics Computer Group, Madison, WI. Other useful codon frequency tables include
35 "Celegans_high.cod", "Celegans_low.cod", "Drosophila_high.cod", "Human_high.cod",

- 28 -

"Maize_high.cod", and "Yeast_high.cod". Other preferred variants are those encoding conservative amino acid changes as described below (e.g., wherein the charge or polarity of the naturally occurring amino acid side chain is not altered substantially by substitution with a different amino acid) as compared to wild type, and/or those designed to either generate a novel glycosylation and/or phosphorylation site(s), or those designed to delete an existing glycosylation and/or phosphorylation site(s).

The gene, cDNA, or fragment thereof encoding a CRP1 or B7RP1 polypeptide can be inserted into an appropriate expression or amplification vector using standard ligation techniques. The vector is typically selected to be functional in the particular host cell employed (i.e., the vector is compatible with the host cell machinery such that amplification of the gene and/or expression of the gene can occur). The gene, cDNA or fragment thereof encoding the CRP1 or B7RP1 polypeptide may be amplified/expressed in prokaryotic, yeast, insect (baculovirus systems) and/or eukaryotic host cells. Selection of the host cell will depend in part on whether the CRP1 or B7RP1 polypeptide or fragment thereof is to be glycosylated and/or phosphorylated. If so, yeast, insect, or mammalian host cells are preferable.

Typically, expression vectors used in any of the host cells will contain sequences for plasmid maintenance and cloning and expression of inserted nucleotide sequences. Such sequences, referred to collectively as "flanking sequences", will include a promoter and other regulatory elements such as an enhancer(s), an origin of replication element, a transcriptional termination element, a complete intron sequence containing a donor and acceptor splice site, a signal peptide sequence, a ribosome binding site

- 29 -

element, a polyadenylation sequence, a polylinker region for inserting the nucleic acid encoding the polypeptide to be expressed, and a selectable marker element. Each of these elements is discussed below.

- 5 Optionally, the vector may contain a "tag" sequence, *i.e.*, an oligonucleotide molecule located at the 5' or 3' end of a CRP1 or B7RP1 polypeptide coding sequence; the oligonucleotide molecule encodes polyHis (such as hexaHis), or other "tag" such as FLAG, HA (hemagglutinin
10 Influenza virus) or *myc* for which commercially available antibodies exist. This tag is typically fused to the polypeptide upon expression of the polypeptide, and can serve as means for affinity purification of a CRP1 or B7RP1 polypeptide from the
15 host cell. Affinity purification can be accomplished, for example, by column chromatography using antibodies against the tag as an affinity matrix. Optionally, the tag can subsequently be removed from a purified CRP1 or B7RP1 polypeptide by various means such as using
20 certain peptidases.

- The human immunoglobulin hinge and Fc region may be fused at either the N-terminus or C-terminus of a CRP1 or B7RP1 polypeptide by one skilled in the art. The subsequent Fc-fusion protein can be purified by use
25 of a Protein A affinity column. An immunoglobulin Fc region is known to exhibit a long pharmacokinetic half-life *in vivo* and proteins fused to an Fc region have been found to exhibit a substantially greater half-life *in vivo* compared to the unfused counterpart.
30 Also, fusion to the Fc region allows for dimerization and/or multimerization of the molecule that may be useful for the bioactivity of some molecules.

- The flanking sequence may be homologous (*i.e.*, from the same species and/or strain as the host
35 cell), heterologous (*i.e.*, from a species other than the host cell species or strain), hybrid (*i.e.*, a

- 30 -

combination of flanking sequences from more than one source), synthetic, or it may be the native CRP1 or B7RP1 nucleic acid flanking sequences. As such, the source of the flanking sequence may be any unicellular
5 prokaryotic or eukaryotic organism, any vertebrate or invertebrate organism, or any plant, provided that the flanking sequence is functional in, and can be activated by, the host cell machinery.

The flanking sequences useful in the vectors
10 of this invention may be obtained by any of several methods well known in the art. Typically, flanking sequences useful herein other than CRP1 or B7RP1 nucleic acid flanking sequence will have been previously identified by mapping and/or by restriction
15 endonuclease digestion and can thus be isolated from the proper tissue source using the appropriate restriction endonucleases. In some cases, the full nucleotide sequence of the flanking sequence may be known. Here, the flanking sequence may be synthesized
20 using the methods described above for nucleic acid synthesis or cloning.

Where all or only a portion of the flanking sequence is known, it may be obtained using PCR and/or by screening a genomic library with suitable
25 oligonucleotide and/or flanking sequence fragments from the same or another species.

Where the flanking sequence is not known, a fragment of DNA containing a flanking sequence may be isolated from a larger piece of DNA that may contain,
30 for example, a coding sequence or even another gene or genes. Isolation may be accomplished by restriction endonuclease digestion using one or more carefully selected enzymes to isolate the proper DNA fragment. After digestion, the desired fragment may be isolated
35 by agarose gel purification, Qiagen® column or other methods known to the skilled artisan. Selection of

- 31 -

suitable enzymes to accomplish this purpose will be readily apparent to one of ordinary skill in the art.

The origin of replication element is typically a part of prokaryotic expression vectors
5 purchased commercially, and aids in the amplification of the vector in a host cell. Amplification of the vector to a certain copy number can, in some cases, be important for optimal expression of the CRP1 or B7RP1 polypeptide. If the vector of choice does not contain
10 an origin of replication site, one may be chemically synthesized based on a known sequence, and ligated into the vector.

The transcription termination element is typically located 3' of the end of a CRP1 or B7RP1
15 polypeptide coding sequence and serves to terminate transcription of a CRP1 or B7RP1 polypeptide. Usually, the transcription termination element in prokaryotic cells is a G-C rich fragment followed by a poly T sequence. While the element is easily cloned from a
20 library or even purchased commercially as part of a vector, it can also be readily synthesized using methods for nucleic acid synthesis such as those described above.

A selectable marker gene element encodes a
25 protein necessary for the survival and growth of a host cell grown in a selective culture medium. Typical selection marker genes encode proteins that (a) confer resistance to antibiotics or other toxins, e.g., ampicillin, tetracycline, or kanamycin for prokaryotic
30 host cells, (b) complement auxotrophic deficiencies of the cell; or (c) supply critical nutrients not available from complex media. Preferred selectable markers are the kanamycin resistance gene, the ampicillin resistance gene, and the tetracycline
35 resistance gene.

- 32 -

The ribosome binding element, commonly called the Shine-Dalgarno sequence (prokaryotes) or the Kozak sequence (eukaryotes), is usually necessary for translational initiation of mRNA. The element is typically located 3' to the promoter and 5' to the coding sequence of the CRP1 or B7RP1 polypeptide to be synthesized. The Shine-Dalgarno sequence is varied but is typically a polypurine (i.e., having a high A-G content). Many Shine-Dalgarno sequences have been identified, each of which can be readily synthesized using methods set forth above and used in a prokaryotic vector.

In those cases where it is desirable for a CRP1 or B7RP1 polypeptide to be secreted from the host cell, a signal sequence may be used to direct export of the polypeptide from the host cell. A CRP1 or B7RP1 transmembrane domain is also inactivated by mutation or deletion to prevent attachment to the host membrane. Typically, the signal sequence is positioned in the coding region of a CRP1 or B7RP1 gene or cDNA, or directly at the 5' end of a CRP1 or B7RP1 gene coding region. Many signal sequences have been identified, and any of them that are functional in the selected host cell may be used in conjunction with a CRP1 or B7RP1 gene or cDNA. Therefore, the signal sequence may be homologous or heterologous to a CRP1 or B7RP1 gene or cDNA, and may be homologous or heterologous to a CRP1 or B7RP1 polypeptides gene or cDNA. Additionally, the signal sequence may be chemically synthesized using methods set forth above.

In most cases, secretion of the polypeptide from the host cell via the presence of a signal peptide will result in the removal of the amino terminal methionine from the polypeptide.

In many cases, transcription of a CRP1 or B7RP1 gene or cDNA is increased by the presence of one

- 33 -

or more introns in the vector; this is particularly true where a CRP1 or B7RP1 polypeptide is produced in eukaryotic host cells, especially mammalian host cells. The introns used may be naturally occurring within a
5 CRP1 or B7RP1 gene, especially where the gene used is a full length genomic sequence or a fragment thereof. Where the intron is not naturally occurring within the gene (as for most cDNAs), the intron(s) may be obtained from another source. The position of the intron with
10 respect to the 5' flanking sequence and a CRP1 or B7RP1 gene is generally important, as the intron must be transcribed to be effective. As such, where a CRP1 or B7RP1 gene inserted into the expression vector is a cDNA molecule, the preferred position for the intron is
15 3' to the transcription start site, and 5' to the polyA transcription termination sequence. Preferably the intron or introns will be located on one side or the other (*i.e.*, 5' or 3') of the cDNA such that it does not interrupt the this coding sequence. Any intron
20 from any source, including any viral, prokaryotic and eukaryotic (plant or animal) organisms, may be used to practice this invention, provided that it is compatible with the host cell(s) into which it is inserted. Also included herein are synthetic introns. Optionally,
25 more than one intron may be used in the vector.

Where one or more of the elements set forth above are not already present in the vector to be used, they may be individually obtained and ligated into the vector. Methods used for obtaining each of the
30 elements are well known to the skilled artisan.

Preferred vectors for practicing this invention are those which are compatible with bacterial, insect, and mammalian host cells. Such vectors include, *inter alia*, pCRII, pCR3, and pcDNA3.1
35 (Invitrogen Company, San Diego, CA), pBSII (Stratagene Company, La Jolla, CA), pET15b (Novagen, Madison, WI),

- 34 -

pGEX (Pharmacia Biotech, Piscataway, NJ), pEGFP-N2 (Clontech, Palo Alto, CA), pETL (BlueBacII; Invitrogen), and pFastBacDual (Gibco/BRL, Grand Island, NY).

5 After the vector has been constructed and a nucleic acid molecule encoding a full length or truncated CRP1 or B7RP1 polypeptide has been inserted into the proper site of the vector, the completed vector may be inserted into a suitable host cell for
10 amplification and/or polypeptide expression.

 Host cells may be prokaryotic host cells (such as *E. coli*) or eukaryotic host cells (such as a yeast cell, an insect cell, or a vertebrate cell). The host cell, when cultured under appropriate conditions,
15 can synthesize a CRP1 or B7RP1 polypeptide which can subsequently be collected from the culture medium (if the host cell secretes it into the medium) or directly from the host cell producing it (if it is not secreted). After collection, a CRP1 or B7RP1
20 polypeptide can be purified using methods such as molecular sieve chromatography, affinity chromatography, and the like.

 Selection of the appropriate host cell for CRP1 or B7RP1 polypeptide production will depend on
25 various factors, such as desired expression levels, polypeptide modifications that are required for activity, such as glycosylation or phosphorylation, or ease of folding into a biologically active molecule.

 Suitable cells or cell lines may be mammalian
30 cells, such as Chinese hamster ovary cells (CHO), human embryonic kidney (HEK) 293 or 293T cells, or 3T3 cells. The selection of suitable mammalian host cells and methods for transformation, culture, amplification, screening and product production and purification are
35 known in the art. Other suitable mammalian cell lines, are the monkey COS-1 and COS-7 cell lines, and the CV-1

- 35 -

cell line. Further exemplary mammalian host cells include primate cell lines and rodent cell lines, including transformed cell lines. Normal diploid cells, cell strains derived from *in vitro* culture of primary tissue, as well as primary explants, are also suitable. Candidate cells may be genotypically deficient in the selection gene, or may contain a dominantly acting selection gene. Other suitable mammalian cell lines include but are not limited to, mouse neuroblastoma N2A cells, HeLa, mouse L-929 cells, 3T3 lines derived from Swiss, Balb-c or NIH mice, BHK or HaK hamster cell lines.

Similarly useful as host cells are bacterial cells. For example, the various strains of *E. coli* (e.g., HB101, DH5 α , DH10, and MC1061) are well-known as host cells in the field of biotechnology. Various strains of *B. subtilis*, *Pseudomonas spp.*, other *Bacillus spp.*, *Streptomyces spp.*, and the like may also be employed in this method.

Many strains of yeast cells known to those skilled in the art are also available as host cells for expression of the polypeptides of the present invention.

Additionally, where desired, insect cell systems may be utilized in the methods of the present invention. Such systems are described for example in Kitts et al. (*Biotechniques*, 14:810-817 (1993)), Lucklow (*Curr. Opin. Biotechnol.*, 4:564-572 (1993)) and Lucklow et al. (*J. Virol.*, 67:4566-4579 (1993)). Preferred insect cells are Sf-9 and Hi5 (Invitrogen, Carlsbad, CA).

"Transformation" or "transfection" of an expression vector into the selected host cell may be accomplished using such methods as calcium chloride, electroporation, microinjection, lipofection or the

- 36 -

DEAE-dextran method. The method selected will in part be a function of the type of host cell to be used. These methods and other suitable methods are well known to the skilled artisan, and are set forth, for example, in Sambrook *et al.*, *supra*.

The host cells transformed or transfected with an expression vector may be cultured using standard media well known to the skilled artisan. The media will usually contain all nutrients necessary for the growth and survival of the cells. Suitable media for culturing *E. coli* cells are, for example, Luria Broth (LB) and/or Terrific Broth (TB). Suitable media for culturing eukaryotic cells are RPMI 1640, MEM, DMEM, all of which may be supplemented with serum and/or growth factors as required by the particular cell line being cultured. A suitable medium for insect cultures is Grace's medium supplemented with yeastolate, lactalbumin hydrolysate, and/or fetal calf serum as necessary.

Typically, an antibiotic or other compound useful for selective growth of the transformed cells only is added as a supplement to the media. The compound to be used will be dictated by the selectable marker element present on the plasmid with which the host cell was transformed. For example, where the selectable marker element is kanamycin resistance, the compound added to the culture medium will be kanamycin.

The amount of CRP1 or B7RP1 polypeptide produced in the host cell can be evaluated using standard methods known in the art. Such methods include, without limitation, Western blot analysis, SDS-polyacrylamide gel electrophoresis, non-denaturing gel electrophoresis, HPLC separation, immunoprecipitation, and/or activity assays such as DNA binding gel shift assays.

- 37 -

If a CRP1 or B7RP1 polypeptide has been designed to be secreted from the host cells, the majority of polypeptide may be found in the cell culture medium. Polypeptides prepared in this way will typically not possess an amino terminal methionine, as it is removed during secretion from the cell. If, however, a CRP1 or B7RP1 polypeptide is not secreted from the host cells, it will be present in the cytoplasm and/or the nucleus (for eukaryotic host cells) or in the cytosol (for gram negative bacteria host cells) and may have an amino terminal methionine.

Purification of a CRP1 or B7RP1 polypeptide from solution can be accomplished using a variety of techniques. If the polypeptide has been synthesized such that it contains a tag such as Hexahistidine (CRP1 or B7RP1/hexaHis) or other small peptide such as FLAG (Eastman Kodak Co., New Haven, CT) or myc (Invitrogen, Carlsbad, CA) at either its carboxyl or amino terminus, it may essentially be purified in a one-step process by passing the tagged polypeptide through an affinity column where the column matrix has a high affinity for the tag or for the polypeptide directly (*i.e.*, a monoclonal antibody specifically recognizing a CRP1 or B7RP1 polypeptide). For example, polyhistidine binds with great affinity and specificity to nickel, thus an affinity column of nickel (such as the Qiagen® nickel columns) can be used for purification of CRP1 or B7RP1/polyHis. (See for example, Ausubel *et al.*, eds., *Current Protocols in Molecular Biology*, Section 10.11.8, John Wiley & Sons, New York (1993)).

Where a CRP1 or B7RP1 polypeptide is prepared without a tag attached, and no antibodies are available, other well known procedures for purification can be used. Such procedures include, without limitation, ion exchange chromatography, molecular sieve chromatography, HPLC, native gel electrophoresis

- 38 -

in combination with gel elution, and preparative isoelectric focusing ("Isoprime" machine/technique, Hoefer Scientific). In some cases, two or more of these techniques may be combined to achieve increased
5 purity.

If it is anticipated that a CRP1 or B7RP1 polypeptide will be found primarily intracellularly, the intracellular material (including inclusion
10 bodies for gram-negative bacteria) can be extracted from the host cell using any standard technique known to the skilled artisan. For example, the host cells can be lysed to release the contents of the periplasm/cytoplasm by French press, homogenization, and/or sonication followed by centrifugation.

15 If a CRP1 or B7RP1 polypeptide has formed inclusion bodies in the cytosol, the inclusion bodies can often bind to the inner and/or outer cellular membranes and thus will be found primarily in the pellet material after centrifugation. The
20 pellet material can then be treated at pH extremes or with chaotropic agent such as a detergent, guanidine, guanidine derivatives, urea, or urea derivatives in the presence of a reducing agent such as dithiothreitol at alkaline pH or tris
25 carboxyethyl phosphine at acid pH to release, break apart, and solubilize the inclusion bodies. The polypeptide in its now soluble form can then be analyzed using gel electrophoresis, immunoprecipitation or the like. If it is desired
30 to isolate a CRP1 or B7RP1 polypeptide, isolation may be accomplished using standard methods such as those set forth below and in Marston *et al.* (*Meth. Enz.*, 182:264-275 (1990)). In some cases, a CRP1 or B7RP1 polypeptide may not be biologically active
35 upon isolation. Various methods for "refolding" or converting the polypeptide to its tertiary structure

- 39 -

and generating disulfide linkages can be used to restore biological activity. Such methods include contacting the solubilized polypeptide with a solution having a pH usually above 7 and in the presence of a particular concentration of an appropriate chaotrope. In most cases the refolding/oxidation solution will also contain a reducing agent or the reducing agent and the corresponding oxidized form in a specific ratio to generate a particular redox potential allowing for disulfide shuffling to occur in the formation of the protein's cysteine bridge(s). Some of the commonly used redox couples include cysteine/cystamine, glutathione (GSH)/dithiobis GSH, cupric chloride, dithiothreitol (DTT)/dithiane DTT, 2-mercaptoethanol (bME)/dithio-b(ME). In many instances a cosolvent is necessary to increase the efficiency of the refolding and the more common reagents used for this purpose include glycerol, polyethylene glycol of various molecular weights, and arginine.

CRP1 or B7RP1 polypeptides, fragments, and/or derivatives thereof may also be prepared by chemical synthesis methods (such as solid phase peptide synthesis) using techniques known in the art such as those set forth by Merrifield *et al.*, (*J. Am. Chem. Soc.*, 85:2149 (1963)), Houghten *et al.* (*Proc Natl Acad. Sci. USA*, 82:5132 (1985)), and Stewart and Young (*Solid Phase Peptide Synthesis*, Pierce Chemical Co., Rockford, IL (1984)). Such polypeptides may be synthesized with or without a methionine on the amino terminus. Chemically synthesized CRP1 or B7RP1 polypeptides or fragments may be oxidized using methods set forth in these references to form disulfide bridges. CRP1 or B7RP1 polypeptides or fragments are expected to have

- 40 -

biological activity comparable to CRP1 or B7RP1 polypeptides produced recombinantly or purified from natural sources, and thus may be used interchangeably with recombinant or natural CRP1 or
5 B7RP1 polypeptide.

Polypeptides

The term " CRP1 or B7RP1 polypeptide" refers to a polypeptide having the amino acid sequence of
10 Figure 1A (SEQ ID NO:2), Figure 2A (SEQ ID NO:7) or Figure 3A (SEQ ID NO:12) and all related polypeptides described herein. Related polypeptides includes allelic variants, splice variants, fragments, derivatives, substitution, deletion, and insertion
15 variants, fusion polypeptides, and orthologs. Such related polypeptides may be mature polypeptides, i.e., polypeptide lacking a signal peptide. A CRP1 or B7RP1 polypeptide may or may not have amino terminal methionine, depending on the manner in which they are
20 prepared.

The term "CRP1 or B7RP1 polypeptide fragment" refers to a peptide or polypeptide that is less than the full length amino acid sequence of a CRP1 or B7RP1 polypeptide as set forth in Figure 1A (SEQ ID NO:2),
25 Figure 2A (SEQ ID NO:7) or Figure 3A (SEQ ID NO:12). Such a fragment may result from truncation at the amino terminus, truncation at the carboxy terminus, and/or a deletion internal to the polypeptide sequence. Such CRP1 or B7RP1 polypeptides fragments may be prepared
30 with or without an amino terminal methionine. In addition, CRP1 or B7RP1 polypeptides fragments may be naturally-occurring splice variants, other splice variants, and fragments resulting from naturally occurring *in vivo* protease activity. Preferred CRP1
35 or B7RP1 polypeptide fragments include soluble forms of CRP1 or B7RP1 which lack a functional transmembrane

- 41 -

domain and comprise part or all of the extracellular domain of either CRP1 or B7RP1.

The term "CRP1 or B7RP1 polypeptide variants" refers to CRP1 or B7RP1 polypeptides whose amino acid sequences contain one or more amino acid sequence substitutions, deletions, and/or additions as compared to the CRP1 or B7RP1 polypeptides amino acid sequences set forth in Figure 1A (SEQ ID NO:2), Figure 2A (SEQ ID NO:7) or Figure 3A (SEQ ID NO:12). Such CRP1 or B7RP1 polypeptides variants can be prepared from the corresponding CRP1 and B7RP1 polypeptides nucleic acid molecule variants, which have a DNA sequence that varies accordingly from the DNA sequences for CRP1 or B7RP1 polypeptides.

As used herein, the term "CRP1 or B7RP1 polypeptide derivatives" refers to CRP1 or B7RP1 polypeptides, variants, or fragments thereof, that have been chemically modified, as for example, by addition of one or more water soluble polymers, N-linked or O-linked carbohydrates, sugars, phosphates, and/or other such molecules, where the molecule or molecules are not naturally attached to wild-type CRP1 or B7RP1 polypeptides. Derivatives further includes deletion of one or more chemical groups naturally attached to the CRP1 or B7RP1 polypeptide.

As used herein, the terms "biologically active CRP1 or B7RP1 polypeptides", "biologically active CRP1 or B7RP1 polypeptide fragments", "biologically active CRP1 or B7RP1 polypeptide variants", and "biologically active CRP1 or B7RP1 polypeptide derivatives" refer to CRP1 or B7RP1 polypeptides having at least one of the activities characteristic of CRP1 or B7RP1. One activity is binding of B7RP1 to CRP1. Another activity is the ability of CRP1 or B7RP1 to stimulate T-cell proliferation and/or activation.

- 42 -

The term "ortholog" refers to a polypeptide that corresponds to a polypeptide identified from a species. For example, mouse and human B7RP1 polypeptides are considered orthologs.

5 The term "mature amino acid sequence" refers to a polypeptide lacking a leader sequence.

 The term "isolated polypeptide" refers to a polypeptide that is free from at least one contaminating polypeptide that is found in its natural
10 environment, and preferably substantially free from any other contaminating mammalian polypeptides.

 The term "identity," as known in the art, is a relationship between the sequences of two or more nucleic acid molecules or two or more polypeptides, as
15 determined by comparing the sequences. In the art, "identity" also means the degree of sequence relatedness between polypeptide or nucleic acid molecule sequences, as the case may be, as determined by the match between strings of nucleotide or amino
20 acid sequences. "Identity" measures the percent of identical matches between two or more sequences with gap alignments addressed by particular computer programs (i.e., "algorithms").

 The term "similarity" refers to a related
25 concept, but in contrast to "identity", a measure of similarity includes both identical matches and conservative substitution matches. Since conservative substitutions apply to polypeptides and not nucleic acid molecules, similarity only deals with polypeptide
30 sequence comparisons. If two polypeptide sequences have, for example, 10/20 identical amino acids, and the remainder are all non-conservative substitutions, then the percent identity and similarity would both be 50%. If in the same example, there are 5 more positions
35 where there are conservative substitutions, then the percent identity remains 50%, but the per cent

- 43 -

similarity would be 75% (15/20). Therefore, in cases where there are conservative substitutions, the degree of similarity between two polypeptide sequences will be higher than the percent identity between those two sequences. "Conservative" amino acid substitutions are described herein below in reference to Table I. Based on Table I, conservative amino acid substitutions are alternate amino acids selected from the same grouping, e.g., basic, acidic, uncharged polar, and non-polar. For example, conservative amino acid substitutions for arginine would be lysine and histidine.

Identity and similarity can be readily calculated by known methods, including but not limited to those described in *Computational Molecular Biology*, Lesk, A.M., ed., Oxford University Press, New York, 1988; *Biocomputing: Informatics and Genome Projects*, Smith, D.W., ed., Academic Press, New York, 1993; *Computer Analysis of Sequence Data, Part 1*, Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey, 1994; *Sequence Analysis in Molecular Biology*, von Heinje, G., Academic Press, 1987; and *Sequence Analysis Primer*, Gribskov, M. and Devereux, J., eds., M. Stockton Press, New York, 1991; and Carillo, H., and Lipman, D., *SIAM J. Applied Math.*, 48:1073 (1988).

Preferred methods to determine identity and/or similarity are designed to give the largest match between the sequences tested. Methods to determine identity and similarity are codified in publicly available computer programs. Preferred computer program methods to determine identity and similarity between two sequences include, but are not limited to, the GCG program package, including GAP (Devereux, J., et al., *Nucleic Acids Research* 12(1):387 (1984); Genetics Computer Group, University of Wisconsin, Madison, WI), BLASTP, BLASTN, and FASTA (Atschul, S.F. et al., *J. Molec. Biol.* 215:403-410

- 44 -

(1990). The BLAST X program is publicly available from the National Center for Biotechnology Information (NCBI) and other sources (*BLAST Manual*, Altschul, S., et al. NCB NLM NIH Bethesda, MD 20894; Altschul, S., et al., *J. Mol. Biol.* 215:403-410 (1990). The well known Smith Waterman algorithm may also be used to determine identity.

By way of example, using the computer algorithm GAP (Genetics Computer Group, University of Wisconsin, Madison, WI), two polypeptides for which the percent sequence identity is to be determined are aligned for optimal matching of their respective amino acids (the "matched span", as determined by the algorithm). A gap opening penalty (which is calculated as 3 X the average diagonal; the "average diagonal" is the average of the diagonal of the comparison matrix being used; the "diagonal" is the score or number assigned to each perfect amino acid match by the particular comparison matrix) and a gap extension penalty (which is usually 1/10 times the gap opening penalty), as well as a comparison matrix such as PAM 250 or BLOSUM 62 are used in conjunction with the algorithm. A standard comparison matrix (see Dayhoff et al., in: *Atlas of Protein Sequence and Structure*, vol. 5, supp.3 (1978) for the PAM250 comparison matrix; see Henikoff et al., *Proc. Natl. Acad. Sci USA*, 89:10915-10919 (1992) for the BLOSUM 62 comparison matrix) is also used by the algorithm.

Preferred parameters for polypeptide sequence comparison include the following:

Algorithm: Needleman and Wunsch, *J. Mol. Biol.* 48:443-453 (1970)

- 45 -

Comparison matrix: BLOSUM 62 from Henikoff and
Henikoff, *Proc. Natl. Acad. Sci. USA* 89:10915-10919
(1992)

Gap Penalty: 12

5 Gap Length Penalty: 4

Threshold of Similarity: 0

The GAP program is useful with the above
parameters. The aforementioned parameters are the
default parameters for polypeptide comparisons (along
10 with no penalty for end gaps) using the GAP algorithm.

Preferred parameters for nucleic acid
molecule sequence comparison include the following:

15 Algorithm: Needleman and Wunsch, *J. Mol Biol.* 48:443-
453 (1970)

Comparison matrix: matches = +10, mismatch = 0

Gap Penalty: 50

Gap Length Penalty: 3

20

The GAP program is also useful with the above
parameters. The aforementioned parameters are the
default parameters for nucleic acid molecule
comparisons.

25

Other exemplary algorithms, gap opening
penalties, gap extension penalties, comparison
matrices, thresholds of similarity, etc. may be used by
those of skill in the art, including those set forth in
30 the Program Manual, Wisconsin Package, Version 9,
September, 1997. The particular choices to be made
will depend on the specific comparison to be made, such
as DNA to DNA, protein to protein, protein to DNA; and
additionally, whether the comparison is between pairs
35 of sequences (in which case GAP or BestFit are
generally preferred) or between one sequence and a

- 46 -

large database of sequences (in which case FASTA or BLASTA are preferred).

Polypeptides that are at least about 70 percent identical will typically have one or more amino acid substitutions, deletions, and/or additions as compared to a wild type CRP1 or B7RP1 polypeptide. In preferred embodiment, polypeptides will have about 75%, 80%, 85%, 90% or 95% identity to CRP1 or B7RP1 polypeptides. Usually, the substitutions of the native residue will be either alanine, or a conservative amino acid so as to have little or no effect on the overall net charge, polarity, or hydrophobicity of the polypeptide. Conservative substitutions are set forth in Table I below.

15

Table I
Conservative Amino Acid Substitutions

Basic:	arginine
	lysine
	histidine
Acidic:	glutamic acid
	aspartic acid
Uncharged Polar:	glutamine
	asparagine
	serine
	threonine
	tyrosine
Non-Polar:	phenylalanine
	tryptophan
	cysteine
	glycine
	alanine
	valine

- 47 -

proline
methionine
leucine
norleucine
isoleucine

CRP1 or B7RP1 polypeptide derivatives are provided by the invention. In one embodiment, 5 chemically modified CRP1 or B7RP1 polypeptide compositions in which CRP1 or B7RP1 polypeptides are linked to a polymer are included within the scope of the present invention. The polymer selected is typically water soluble so that the protein to which 10 it is attached does not precipitate in an aqueous environment, such as a physiological environment. The polymer selected is usually modified to have a single reactive group, such as an active ester for acylation or an aldehyde for alkylation, so that the 15 degree of polymerization may be controlled as provided for in the present methods. The polymer may be of any molecular weight, and may be branched or unbranched. Included within the scope of the invention is a mixture of polymers. Preferably, for 20 therapeutic use of the end-product preparation, the polymer will be pharmaceutically acceptable.

The water soluble polymer or mixture thereof may be selected from the group consisting of, for example, polyethylene glycol (PEG), 25 monomethoxy-polyethylene glycol, dextran, cellulose, or other carbohydrate based polymers, poly-(N-vinyl pyrrolidone) polyethylene glycol, propylene glycol homopolymers, a polypropylene oxide/ethylene oxide co-polymer, polyoxyethylated polyols (e.g., 30 glycerol) and polyvinyl alcohol.

- 48 -

For the acylation reactions, the polymer(s) selected should have a single reactive ester group. For reductive alkylation, the polymer(s) selected should have a single reactive aldehyde group. A preferred reactive aldehyde is polyethylene glycol propionaldehyde, which is water stable, or mono C1-C10 alkoxy or aryloxy derivatives thereof (see U.S. Patent No. 5,252,714).

Pegylation of CRP1 or B7RP1 polypeptides may be carried out by any of the pegylation reactions known in the art, as described for example in the following references: *Focus on Growth Factors* 3: 4-10 (1992); EP 0 154 316; and EP 0 401 384. Preferably, the pegylation is carried out via an acylation reaction or an alkylation reaction with a reactive polyethylene glycol molecule (or an analogous reactive water-soluble polymer) as described below.

A particularly preferred water-soluble polymer for use herein is polyethylene glycol, abbreviated PEG. As used herein, polyethylene glycol is meant to encompass any of the forms of PEG that have been used to derivatize other proteins, such as mono-(C1-C10) alkoxy- or aryloxy-polyethylene glycol.

In general, chemical derivatization may be performed under any suitable conditions used to react a biologically active substance with an activated polymer molecule. Methods for preparing pegylated CRP1 and B7RP1 polypeptides will generally comprise the steps of (a) reacting the polypeptide with polyethylene glycol (such as a reactive ester or aldehyde derivative of PEG) under conditions whereby CRP1 or B7RP1 polypeptide becomes attached to one or more PEG groups, and (b) obtaining the reaction product(s). In general, the optimal reaction conditions for the acylation reactions will be determined based on known parameters and the desired result. For example, the larger the ratio of

- 49 -

PEG: protein, the greater the percentage of poly-pegylated product.

Generally, conditions which may be alleviated or modulated by administration of CRP1 or B7RP1 polymer
5 conjugates include those described herein for non-conjugated CRP1 or B7RP1 polypeptides. However, the conjugated disclosed herein may have additional activities, enhanced or reduced biological activity, or other characteristics, such as increased or decreased
10 half-life, as compared to the non-derivatized molecules.

CRP1 or B7RP1 polypeptides, fragments variants, and derivatives, may be employed alone, together, or in combination with other pharmaceutical
15 compositions. CRP1 or B7RP1 polypeptides, fragments, variants, and derivatives may be used in combination with cytokines, growth factors, antibiotics, anti-inflammatories, and/or chemotherapeutic agents as is appropriate for the indication being treated.

20

The invention provides for selective binding agents of CRP1 or B7RP1. A selective binding agent refers to a molecule having specificity for CRP1 or B7RP1 and may include a protein, peptide, nucleic acid,
25 carbohydrate, lipid or small molecular weight compound. A selective binding agent interacts either with CRP1 or B7RP1 and in turn regulates the binding of CRP1 to B7RP1. In one embodiment, a selective binding agent partially or completely blocks the binding of CRP1 to
30 B7RP1 and partially or completely inhibits at least one biological activity of CRP-1 or B7RP-1, such as immune costimulatory activity. In another embodiment, the selective binding agent is an antibody. The antibody may be immunoreactive with either CRP1 or B7RP1 and is
35 preferably immunoreactive with B7RP1. In yet another embodiment of the invention, an antibody reactive with

- 50 -

B7RP1 binds to an eptiope on B7RP1 such that binding to CRP1 is partially or completely blocked and at least one biological activity of B7RP1, such as immune costimulatory activity, is partially or completely inhibited. The term partially inhibited means that at least a detectable level of inhibition has occurred. The term completely inhibited means that no further increase in inhibition has occurred.

CRP1 or B7RP1 polypeptides, fragments, variants, and/or derivatives may be used to prepare antibodies using methods known in the art. Thus, antibodies that react with the CRP1 or B7RP1 polypeptides, as well as reactive fragments of such antibodies, are also contemplated as within the scope of the present invention. The antibodies may be polyclonal, monoclonal, recombinant, chimeric, single-chain and/or bispecific. Typically, the antibody or fragment thereof will either be of human origin, or will be "humanized", i.e., prepared so as to prevent or minimize an immune reaction to the antibody when administered to a patient. The antibody fragment may be any fragment that is reactive with CRP1 and B7RP1 polypeptides of the present invention, such as, Fab, Fab', etc. Also provided by this invention are the hybridomas generated by presenting any CRP1 or B7RP1 polypeptide or fragments thereof as an antigen to a selected mammal, followed by fusing cells (e.g., spleen cells) of the mammal with certain cancer cells to create immortalized cell lines by known techniques. The methods employed to generate such cell lines and antibodies directed against all or portions of a human CRP1 or B7RP1 polypeptide of the present invention are also encompassed by this invention.

Monoclonal antibodies of the invention include "chimeric" antibodies in which a portion of the

- 51 -

heavy and/or light chain is identical with or homologous to corresponding sequence in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chains(s) is identical with or homologous to corresponding sequence in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such antibodies, so long as they exhibit the desired biological activity (U.S. Patent No. 4,816,567; Morrison, et al., Proc. Natl. Acad. Sci. 81, 6851-6855 (1985)).

In a preferred embodiment, the chimeric anti-CRP1 or B7RP1 antibody is a "humanized" antibody. Methods for humanizing non-human antibodies are well known in the art. Generally, a humanized antibody has one or more amino acid residues introduced into it from a source which is non-human. Humanization can be performed following methods known in the art (Jones, et al., Nature 321, 522-525 (1986); Riechmann, et al., Nature, 332, 323-327 (1988); Verhoeyen, et al., Science 239, 1534-1536 (1988)), by substituting rodent complementarily-determining regions (CDRs) for the corresponding regions of a human antibody.

Also encompassed by the invention are fully human anti-CRP1 or anti-B7RP1 antibodies. Such antibodies may be produced by immunization with a CRP1 or B7RP1 antigen of transgenic animals (e.g., mice) that are capable of producing a repertoire of human antibodies in the absence of endogenous immunoglobulin production. See, for example, Jakobovits, et al., Proc. Natl. Acad. Sci. 90, 2551-2555 (1993); Jakobovits, et al., Nature 362, 255-258 (1993). Human antibodies can also be produced in phage-display libraries (Hoogenboom, et al., J. Mol. Biol. 227, 381 (1991); Marks, et al., J. Mol. Biol. 222, 581 (1991)).

- 52 -

Selective binding agents of the invention may be used to regulate the binding of CRP1 to B7RP1 and regulate at least one biological activity mediated by CRP1 and B7RP1 such as immune co-stimulation. An
5 example of such selective binding agents are antibodies immunoreactive with either CRP1 or B7RP1. The antibodies may be used therapeutically, such as to inhibit binding of the CRP1 and B7RP1 polypeptide to its binding partner. The antibodies may further be
10 used for *in vivo* and *in vitro* diagnostic purposes, such as in labeled form to detect the presence of CRP1 and B7RP1 polypeptide in a body fluid or cell sample.

Pharmaceutical Compositions and Administration

15 Pharmaceutical compositions of CRP1 or B7RP1 polypeptides are within the scope of the present invention. Such compositions may comprise a therapeutically effective amount of the polypeptide or fragments, variants, or derivatives in admixture with a
20 pharmaceutically acceptable carrier. In preferred embodiments, pharmaceutical compositions comprise CRP1 or B7RP1 polypeptides as soluble forms which comprise part or all of a CRP1 or B7RP1 extracellular domain. Typically, a CRP1 and B7RP1 polypeptide therapeutic
25 compound will be administered in the form of a composition comprising purified polypeptide, fragment, variant, or derivative in conjunction with one or more physiologically acceptable carriers, excipients, or diluents. The carrier material may be water for
30 injection, preferably supplemented with other materials common in solutions for administration to mammals. Neutral buffered saline or saline mixed with serum albumin are exemplary appropriate carriers. Preferably, the product is formulated as a lyophilizate
35 using appropriate excipients (e.g., sucrose). Other standard carriers, diluents, and excipients may be

- 53 -

included as desired. Other exemplary compositions comprise Tris buffer of about pH 7.0-8.5, or acetate buffer of about pH 4.0-5.5, which may further include sorbitol or a suitable substitute therefor.

5 CRP1 or B7RP1 pharmaceutical compositions can be administered parenterally. Alternatively, the compositions may be administered intravenously or subcutaneously. When systemically administered, the therapeutic compositions for use in this invention may
10 be in the form of a pyrogen-free, parenterally acceptable aqueous solution. The preparation of such pharmaceutically acceptable protein solutions, with due regard to pH, isotonicity, stability and the like, is within the skill of the art.

15 Therapeutic formulations of CRP1 and B7RP1 polypeptide compositions useful for practicing the present invention may be prepared for storage by mixing the selected composition having the desired degree of purity with optional physiologically acceptable
20 carriers, excipients, or stabilizers (*Remington's Pharmaceutical Sciences*, 18th Edition, A.R. Gennaro, ed., Mack Publishing Company (1990)) in the form of a lyophilized cake or an aqueous solution. Acceptable carriers, excipients or stabilizers are nontoxic to
25 recipients and are preferably inert at the dosages and concentrations employed, and include buffers such as phosphate, citrate, or other organic acids; antioxidants such as ascorbic acid; low molecular weight polypeptides; proteins, such as serum albumin,
30 gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, arginine or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating
35 agents such as EDTA; sugar alcohols such as mannitol or sorbitol; salt-forming counterions such as sodium;

- 54 -

and/or nonionic surfactants such as Tween, pluronics or polyethylene glycol (PEG).

An effective amount of a CRP1 or B7RP1 polypeptide composition(s) to be employed therapeutically will depend, for example, upon the therapeutic objectives such as the indication for which the CRP1 and B7RP1 polypeptide is being used, the route of administration, and the condition of the patient. Accordingly, it will be necessary for the therapist to titer the dosage and modify the route of administration as required to obtain the optimal therapeutic effect. A typical daily dosage may range from about 0.1 µg/kg to up to 100 mg/kg or more, depending on the factors mentioned above. Typically, a clinician will administer the composition until a dosage is reached that achieves the desired effect. The composition may therefore be administered as a single dose, or as two or more doses (which may or may not contain the same amount of a CRP1 or B7RP1 polypeptide) over time, or as a continuous infusion via implantation device or catheter.

As further studies are conducted, information will emerge regarding appropriate dosage levels for treatment of various conditions in various patients, and the ordinary skilled worker, considering the therapeutic context, the type of disorder under treatment, the age and general health of the recipient, will be able to ascertain proper dosing.

The CRP1 or B7RP1 polypeptide composition to be used for *in vivo* administration must be sterile. This is readily accomplished by filtration through sterile filtration membranes. Where the composition is lyophilized, sterilization using these methods may be conducted either prior to, or following, lyophilization and reconstitution. The composition for parenteral

- 55 -

administration ordinarily will be stored in lyophilized form or in solution.

Therapeutic compositions generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle.

The route of administration of the composition is in accord with known methods, e.g. oral, injection or infusion by intravenous, intraperitoneal, intracerebral (intraparenchymal), intracerebroventricular, intramuscular, intraocular, intraarterial, or intralesional routes, or by sustained release systems or implantation device which may optionally involve the use of a catheter. Where desired, the compositions may be administered continuously by infusion, bolus injection or by implantation device.

Alternatively or additionally, the composition may be administered locally via implantation into the affected area of a membrane, sponge, or other appropriate material on to which CRP1 and B7RP1 polypeptide has been absorbed.

Where an implantation device is used, the device may be implanted into any suitable tissue or organ, and delivery of a CRP1 or B7RP1 polypeptide may be directly through the device via bolus, or via continuous administration, or via catheter using continuous infusion. A CRP1 or B7RP1 polypeptide may be administered in a sustained release formulation or preparation. Suitable examples of sustained-release preparations include semipermeable polymer matrices in the form of shaped articles, e.g. films, or microcapsules. Sustained release matrices include polyesters, hydrogels, polylactides (U.S. 3,773,919, EP 58,481), copolymers of L-glutamic acid and gamma ethyl-L-glutamate (Sidman et al, *Biopolymers*, 22: 547-556

- 56 -

(1983)), poly (2-hydroxyethyl-methacrylate) (Langer et al., *J. Biomed. Mater. Res.*, 15: 167-277 (1981)] and Langer, *Chem. Tech.*, 12: 98-105 (1982)), ethylene vinyl acetate (Langer et al., *supra*) or poly-D(-)-3-hydroxybutyric acid (EP 133,988). Sustained-release compositions also may include liposomes, which can be prepared by any of several methods known in the art (e.g., Eppstein et al., *Proc. Natl. Acad. Sci. USA*, 82: 3688-3692 (1985); EP 36,676; EP 88,046; EP 143,949).

10

In some cases, it may be desirable to use CRP1 or B7RP1 polypeptide compositions in an *ex vivo* manner. Here, cells, tissues, or organs that have been removed from the patient are exposed to a CRP1 or B7RP1 polypeptide compositions after which the cells, tissues and/or organs are subsequently implanted back into the patient.

In other cases, a CRP1 or B7RP1 polypeptide may be delivered through implanting into patients certain cells that have been genetically engineered, using methods such as those described herein, to express and secrete the polypeptides, fragments, variants, or derivatives. Such cells may be animal or human cells, and may be derived from the patient's own tissue or from another source, either human or non-human. Optionally, the cells may be immortalized. However, in order to decrease the chance of an immunological response, it is preferred that the cells be encapsulated to avoid infiltration of surrounding tissues. The encapsulation materials are typically biocompatible, semi-permeable polymeric enclosures or membranes that allow release of the protein product(s) but prevent destruction of the cells by the patient's immune system or by other detrimental factors from the surrounding tissues.

- 57 -

Methods used for membrane encapsulation of cells are familiar to the skilled artisan, and preparation of encapsulated cells and their implantation in patients may be accomplished without undue experimentation. See, e.g., U.S. Patent Nos. 4,892,538; 5,011,472; and 5,106,627. A system for encapsulating living cells is described in PCT WO 91/10425 (Aebischer et al.). Techniques for formulating a variety of other sustained or controlled delivery means, such as liposome carriers, bio-erodible particles or beads, are also known to those in the art, and are described, for example, in U.S. Patent No. 5,653,975. The cells, with or without encapsulation, may be implanted into suitable body tissues or organs of the patient.

As discussed above, it may be desirable to treat cell preparations with one or more CRP1 or B7RP1 polypeptides, variants, derivatives and/or fragments. This can be accomplished by exposing, for example, cells comprising T-cells, such as bone marrow cells, to the polypeptide, variant, derivative, or fragment directly, where it is in a form that is permeable to the cell membrane. For example, cells comprising T-cells may be exposed to a B7RP1 polypeptide in order to activate T-cell function and the cells so treated are implanted in the patient.

Alternatively, gene therapy can be employed. One manner in which gene therapy can be applied is to use a CRP1 or B7RP1 gene (either genomic DNA, cDNA, and/or synthetic DNA encoding a CRP1 or B7RP1 polypeptide, or a fragment, variant, or derivative thereof) which may be operably linked to a constitutive or inducible promoter to form a "gene therapy DNA construct". The promoter may be homologous or heterologous to the endogenous CRP1 or B7RP1 gene, provided that it is active in the cell or tissue type

- 58 -

into which the construct will be inserted. Other components of the gene therapy DNA construct may optionally include, as required, DNA molecules designed for site-specific integration (e.g., endogenous
5 flanking sequences useful for homologous recombination), tissue-specific promoter, enhancer(s) or silencer(s), DNA molecules capable of providing a selective advantage over the parent cell, DNA molecules useful as labels to identify transformed cells,
10 negative selection systems, cell specific binding agents (as, for example, for cell targeting) cell-specific internalization factors, and transcription factors to enhance expression by a vector as well as factors to enable vector manufacture.

15 This gene therapy DNA construct can then be introduced into the patient's cells (either *ex vivo* or *in vivo*). One means for introducing the gene therapy DNA construct is via viral vectors. Suitable viral vectors typically used in gene therapy for delivery of
20 gene therapy DNA constructs include, without limitation, adenovirus, adeno-associated virus, herpes simplex virus, lentivirus, papilloma virus, and retrovirus vectors. Some of these vectors, such as retroviral vectors, will deliver the gene therapy DNA
25 construct to the chromosomal DNA of the patient's cells, and the gene therapy DNA construct can integrate into the chromosomal DNA; other vectors will function as episomes and the gene therapy DNA construct will remain in the cytoplasm. The use of gene therapy
30 vectors is described, for example, in U.S. Patent Nos. 5,672,344, 5,399,346.

Alternative means to deliver gene therapy DNA constructs to a patient's cells without the use of viral vectors include, without limitation, liposome-mediated transfer, direct injection of naked DNA,
35 receptor-mediated transfer (ligand-DNA complex),

- 59 -

electroporation, calcium phosphate precipitation, and microparticle bombardment (e.g., "gene gun"). See U.S. Patent Nos. 4,970,154, WO 96/40958, 5,679,559, 5,676,954, and 5,593,875.

5 Another means to increase endogenous CRP1 or B7RP1 polypeptide expression in a cell via gene therapy is to insert one or more enhancer elements into the CRP1 or B7RP1 polypeptide promoter, where the enhancer element(s) can serve to increase transcriptional
10 activity of a CRP1 or B7RP1 polypeptide gene. The enhancer element(s) used will be selected based on the tissue in which one desires to activate the gene(s); enhancer elements known to confer promoter activation in that tissue will be selected. For example, if a
15 CRP1 or B7RP1 polypeptide is to be "turned on" in T-cells, the *lck* promoter enhancer element may be used. Here, the functional portion of the transcriptional element to be added may be inserted into a fragment of DNA containing a CRP1 or B7RP1 polypeptide promoter
20 (and optionally, vector, 5' and/or 3' flanking sequence, etc.) using standard cloning techniques. This construct, known as a "homologous recombination construct" can then be introduced into the desired cells either *ex vivo* or *in vivo*.

25 Gene therapy can be used to decrease CRP1 or B7RP1 polypeptide expression by modifying the nucleotide sequence of the endogenous promoter(s). Such modification is typically accomplished via homologous recombination methods. For example, a DNA
30 molecule containing all or a portion of the promoter of a CRP1 or B7RP1 gene(s) selected for inactivation can be engineered to remove and/or replace pieces of the promoter that regulate transcription. Here, the TATA box and/or the binding site of a transcriptional
35 activator of the promoter may be deleted using standard molecular biology techniques; such deletion can inhibit

- 60 -

promoter activity thereby repressing transcription of the corresponding CRP1 or B7RP1 gene. Deletion of the TATA box or transcription activator binding site in the promoter may be accomplished by generating a DNA
5 construct comprising all or the relevant portion of a CRP1 or B7RP1 polypeptide promoter(s) (from the same or a related species as a CRP1 or B7RP1 gene(s) to be regulated) in which one or more of the TATA box and/or transcriptional activator binding site nucleotides are
10 mutated via substitution, deletion and/or insertion of one or more nucleotides such that the TATA box and/or activator binding site has decreased activity or is rendered completely inactive. This construct, which also will typically contain at least about 500 bases of
15 DNA that correspond to the native (endogenous) 5' and 3' flanking regions of the promoter segment that has been modified, may be introduced into the appropriate cells (either *ex vivo* or *in vivo*) either directly or via a viral vector as described above. Typically,
20 integration of the construct into the genomic DNA of the cells will be via homologous recombination, where the 5' and 3' flanking DNA sequences in the promoter construct can serve to help integrate the modified promoter region via hybridization to the endogenous
25 chromosomal DNA.

Other gene therapy methods may also be employed where it is desirable to inhibit one or more CRP1 or B7RP1 polypeptides. For example, antisense DNA or RNA molecules, which have a sequence that is
30 complementary to at least a portion of a selected CRP1 or B7RP1 polypeptide gene(s) can be introduced into the cell. Typically, each such antisense molecule will be complementary to the start site (5' end) of each selected CRP1 or B7RP1 gene. When the antisense
35 molecule then hybridizes to the corresponding CRP1 or

- 61 -

B7RP1 polypeptide mRNA, translation of this mRNA is prevented.

Alternatively, gene therapy may be employed to create a dominant-negative inhibitor of one or more CRP1 or B7RP1 polypeptides. In this situation, the DNA encoding a mutant full length or truncated polypeptide of each selected CRP1 or B7RP1 polypeptide can be prepared and introduced into the cells of a patient using either viral or non-viral methods as described above. Each such mutant is typically designed to compete with endogenous polypeptide in its biological role.

Agonists and Antagonists

The invention also provides for agonists and antagonists of CRP1 or B7RP1 which regulate the activity of either or both molecules. Agonists and antagonists may be identified from test molecules which alter the binding of B7RP1 to CRP1.

The term "test molecule(s)" refers to the molecule(s) that is/are under evaluation for the ability to bind a CRP1 or B7RP1 polypeptide and thereby alter the binding of B7RP1 to CRP1. Preferably, the test molecule will bind with an affinity constant of at least about 10^6 M.

A variety of assays may be used to measure binding of B7RP1 to CRP1. These assays may be used to screen test molecules for their ability to increase or decrease the rate or extent of binding of B7RP1 to CRP1. In one type of assay, a CRP1 polypeptide, preferably a soluble form of CRP1 such as an extracellular domain, is immobilized by attachment to the bottom of the wells of a microtiter plate. Radiolabeled B7RP1 and the test molecule(s) can then be added either one at a time (in either order) or simultaneously to the wells. After incubation, the

- 62 -

wells can be washed and counted using a scintillation counter for radioactivity to determine the extent of binding to CRP1 protein by B7RP1. Typically, the molecules will be tested over a range of concentrations, and a series of control wells lacking one or more elements of the test assays can be used for accuracy in evaluation of the results. An alternative to this method involves reversing the "positions" of the proteins, i.e., immobilizing B7RP1 to the microtiter plate wells, incubating with the test molecule and radiolabeled CRP1, and determining the extent of CRP1 binding (see, for example, chapter 18 of *Current Protocols in Molecular Biology*, Ausubel et al., eds., John Wiley & Sons, New York, NY [1995]).

As an alternative to radiolabelling, CRP1 or B7RP1 may be conjugated to biotin and the presence of biotinylated protein can then be detected using streptavidin linked to an enzyme, such as horse radish peroxidase [HRP] or alkaline phosphatase [AP], that can be detected colorometrically, or by fluorescent tagging of streptavidin. An antibody directed to CRP1 or B7RP1 that is conjugated to biotin may also be used and can be detected after incubation with enzyme-linked streptavidin linked to AP or HRP

CRP1 and B7RP1 may also be immobilized by attachment to agarose beads, acrylic beads or other types of such inert substrates. The substrate-protein complex can be placed in a solution containing the complementary protein and the test compound; after incubation, the beads can be precipitated by centrifugation, and the amount of binding between CRP1 and B7RP1 can be assessed using the methods described above. Alternatively, the substrate-protein complex can be immobilized in a column and the test molecule and complementary protein passed over the column. Formation of a complex between CRP1 and B7RP1 can then

- 63 -

be assessed using any of the techniques set forth above, *i.e.*, radiolabelling, antibody binding, or the like.

Another type of in vitro assay that is useful
5 for identifying a test molecule which increases or decreases formation of an CRP1/B7RP1 complex is a surface plasmon resonance detector system such as the Biacore assay system (Pharmacia, Piscataway, NJ). The Biacore system may be carried out using the
10 manufacturer's protocol. This assay essentially involves covalent binding of either CRP1 or B7RP1 to a dextran-coated sensor chip which is located in a detector. The test compound and the other complementary protein can then be injected into the
15 chamber containing the sensor chip either simultaneously or sequentially and the amount of complementary protein that binds can be assessed based on the change in molecular mass which is physically associated with the dextran-coated side of the of the
20 sensor chip; the change in molecular mass can be measured by the detector system.

In some cases, it may be desirable to evaluate two or more test compounds together for use in increasing or decreasing formation of a CRP1/B7RP1
25 complex. In these cases, the assays set forth above can be readily modified by adding such additional test compound(s) either simultaneously with, or subsequently to, the first test compound. The remainder of steps in the assay are as set forth above.

30 In vitro assays such as those described above may be used advantageously to screen rapidly large numbers of compounds for effects on complex formation by CRP1 and B7RP1. The assays may be automated to screen compounds generated in phage display, synthetic
35 peptide and chemical synthesis libraries.

- 64 -

Compounds which increase or decrease complex formation of CRP1 and B7RP1 may also be screened in cell culture using cells and cell lines expressing either polypeptide. Cells and cell lines may be
5 obtained from any mammal, but preferably will be from human or other primate, canine, or rodent sources. The binding of B7RP1 to cells expressing CRP1 on the surface is evaluated in the presence or absence of test molecules and the extent of binding may be determined
10 by, for example, flow cytometry using a biotinylated antibody to B7RP1. Cell culture assays may be used advantageously to further evaluate compounds that score positive in protein binding assays described above.

15 Therapeutic Uses

Polypeptides of the invention, and agonists and antagonists thereof, may be used to regulate T-cell function. Agonists and antagonists include those molecules which regulate CRP1 and/or B7RP1 activity and
20 either increase or decrease at least one activity of a CRP1 or B7RP1 protein such as one activity associated with T-cell functions, for example, T-cell activation. Agonists or antagonists may be co-factors, such as a protein, peptide, carbohydrate, lipid, or small
25 molecular weight molecule, which interact with either CRP1 or B7RP1 and thereby regulate their activity. Potential polypeptide agonists or antagonists include antibodies that react with either soluble or membrane-bound forms of CRP1 or B7RP1 which comprise part or all
30 of the extracellular domains of the said proteins. Molecules that regulate CRP1 or B7RP1 expression typically include nucleic acids encoding CRP1 or B7RP1 protein that can act as anti-sense regulators of expression.

35 CRP1 or B7RP1 polypeptides, and agonists and antagonists thereof, may be used in the treatment of

- 65 -

autoimmune disease, graft survival, immune cell activation for inhibiting tumor cell growth, T-cell dependent B-cell mediated diseases, and cancer gene immunotherapy. In one embodiment, antagonists or
5 inhibitors of CRP1 and/or B7RP1 function may be beneficial to alleviate symptoms in diseases with chronic immune cell dysfunction. Autoimmune diseases, such as systemic lupus erythematosus, rheumatoid arthritis, immune thrombocytopenic purpura (ITP), and
10 psoriasis, may be treated with antagonists or inhibitors of CRP-1/B7RP-1. In addition, chronic inflammatory diseases, such as inflammatory bowel disease (Crohn's disease and ulcerative colitis), Grave's disease, Hashimoto's thyroiditis, and diabetes
15 mellitis, may also be treated with inhibitors to CRP-1/B7RP-1. As described in Example 18, CRP-1-Fc inhibits and B7RP-1-Fc enhances, the onset of disease in a rodent rheumatoid arthritis disease model. These opposite effects in this model support an agonistic
20 role for the B7RP-1-Fc protein and an antagonistic role for the CRP-1-Fc protein. The results also illustrate how T-cell responses can be regulated by manipulation of this pathway and the significance of this pathway in the progression of rheumatoid arthritis. In addition,
25 as described in Example 19, expression of B7RP-1-Fc *in vivo* stimulates an inflammatory bowel disease (IBD) phenotype in transgenic mice. This example supports the role for B7RP-1/CRP-1 in the development of inflammation in the intestine. Therefore, antagonists
30 of the B7RP-1/CRP-1 pathway may be used to treat human IBD.

Antagonists of CRP1 or B7RP1 may be used as immunosuppressive agents for bone marrow and organ transplantation and may be used to prolong graft
35 survival. Such antagonists may provide significant advantages over existing treatment. Bone marrow and

- 66 -

organ transplantation therapy must contend with T-cell mediated rejection of the foreign cells or tissue by the host. Present therapeutic regimens for inhibiting T-cell mediated rejection involve treatment with the
5 drugs cyclosporine or FK506. While drugs are effective, patients suffer from serious side effects, including hepatotoxicity, nephrotoxicity, and neurotoxicity. The target for the cyclosporin/FK506 class of therapeutics is calcineurin, a phosphatase
10 with ubiquitous expression. Since CRP1 expression is restricted to T-cells, inhibitors of CRP1 or B7RP1 may lack the severe side effects observed with the use of the present immunotherapeutic agents.

Antagonists of CRP1 or B7RP1 may be used as
15 immunosuppressive agents for autoimmune disorders, such as rheumatoid arthritis, psoriasis, multiple sclerosis, diabetes, and systemic lupus erythematosus.

Antagonists of the CRP1/B7RP1-mediated costimulatory pathway may also be used to alleviate
20 toxic shock syndrome, inflammatory bowel disease, allosensitization due to blood transfusions, T-cell dependent B-cell mediated diseases, and the treatment of graft vs. host disease.

Antibodies, soluble proteins comprising for
25 example extracellular domains, and other regulators of CRP1 or B7RP1 that result in prolonged or enhanced T-cell activation can be used to increased the immune response to tumors. Example 20 shows B7RP-1-Fc can inhibit tumor cell growth in mice. Similarly, human
30 B7RP-1-Fc, or other activators of the B7RP-1/CRP-1 pathway, may be used to enhance immune responses against human tumors. Anti-tumor activity is generally considered to have a strong cytolytic T-lymphocyte component. In fact, the anti-tumor effects of B7-Fc
35 fusion proteins (Sturmhoefel et al., Cancer Res. 59: 4964-4972, 1999) were mediated by cytolytic CD8+ T-

- 67 -

cells. Since CRP-1 is also expressed on cytolytic CD8+ T-cells (Example 9), it is probable that the anti-tumor effects demonstrated in Example 20 were due to B7RP-1-Fc action on CD8+ cells. The B7RP-1/CRP-1 pathway can
5 also be manipulated to regulate CTL response in a number of other clinical settings, including allograft transplantation, graft vs. host disease, and autoimmune diseases.

Gene therapy using B7RP1 genes of the
10 invention may be used in cancer immunotherapy. B7RP1 genes introduced into cancer cells can transform them into antigen presenting cells that can be recognized by the T-cells of the immune system when introduced back into an animal. Recognition of the transfected tumor
15 cells by the T-cells results in eradication of both tumors cells expressing, or not expressing, the B7RP1 gene. This immunotherapy approach may be used for various leukemias, sarcomas, melanomas, adenocarcinomas, breast carcinomas, prostate tumors,
20 lung carcinomas, colon carcinomas and other tumors. This invention encompasses using the B7RP1 gene in a similar manner to enhance T-cell activation in response to variety of tumors.

As described in Example 14, the phenotype of
25 transgenic mice expressing B7RP1 indicates that B7RP1 is important in the control of antibody production. Agonists and antagonists of B7RP1 protein activity may be useful in therapeutic indications that call for the inhibition or enhancement of antibody production.

30 For instance, many vaccines act by eliciting an effective and specific antibody response. Some vaccines, especially those against intestinal micro-organisms (e.g. Hepatitis A virus, and Salmonellas), elicit a short-lived antibody response. It is desirable
35 to potentiate and prolong this response in order to increase the effectiveness of the vaccine. Therefore,

- 68 -

soluble B7RP1 or activating antibodies to CRP1 may serve as a vaccine adjuvant.

Anti-viral responses may also be enhanced by activators or agonists of the B7RP-1/CRP-1 pathway. The data in Example 20 indicate that cellular immunity is enhanced by B7RP-1-Fc. The enhancement of cellular immune functions by B7RP-1-Fc, or other activators of the B7RP-1/CRP-1 pathway, may also be beneficial in eliminating virus-infected cells. In a complementary fashion, B7RP-1-Fc has effects on humoral immune functions that may enhance antibody mediated responses as observed in Example 13 that may function to help clear free-virus from the body.

Conversely, there are a number of clinical conditions that would be ameliorated by the inhibition of antibody production. Hypersensitivity is a normally beneficial immune response that is exaggerated or inappropriate, and leads to inflammatory reactions and tissue damage. Hypersensitivity reactions which are antibody-mediated may be particularly susceptible to antagonism by inhibitors of B7RP1 activity. Allergies, hay fever, asthma, and acute edema cause type I hypersensitivity reactions, and these reactions may be suppressed by protein, antibody or small molecule inhibitors of B7RP1 activity.

Diseases that cause antibody-mediated hypersensitivity reactions, including systemic lupus erythematosus, arthritis (rheumatoid arthritis, reactive arthritis, psoriatic arthritis), nephropathies (glomerulo-nephritis, membranous, mesangiocapillary, focal segmental, focal necrotizing, crescentic, proliferative - tubulopathies), skin disorders (pemphigus and pemphigoid, erythema nodosum), endocrinopathies (thyroiditis - Grave's, Hashimoto's - insulin dependent diabetes mellitus), various pneumopathies (especially extrinsic alveolitis),

- 69 -

various vasculopathies, coeliac disease, with aberrant production of IgA, many anemias and thrombocytopenias, Guillain-Barre Syndrome, and myasthenia gravis, may be treated with B7RP1 antagonists.

5 In addition, lymphoproliferative disorders, such as multiple myeloma, Waldenstrom's macroglobulinemia, and crioglobulinemias, may be inhibited by protein, antibody, or small molecule antagonists of B7RP1.

10 Finally, graft versus host disease, an "artificial" immune disorder, may benefit from the inhibition of antibody production by B7RP1 antagonists.

 The following examples are offered to more
15 fully illustrate the invention, but are not construed as limiting the scope thereof.

Example 1

20 CRP1 cDNA and Amino Acid Sequence

 Female C57/ Black 6 mice were sacrificed, and the small intestines were excised, and the Peyer's patches were removed. The small intestine tissue was
25 sliced open and washed to remove mucus and other debris. The epithelial layer, which contains the intestinal intraepithelial cells (iIELs), was released by gentle agitation in RPMI-1640 supplemented with 1 mM dithiothreitol (DTT), for 20 minutes at 37°C.
30 Disassociated cells were passed through a 100 μ filter, washed in 50 ml of RPMI-1640, mixed to further break up clumps of cells, and then passed through a 40 μ strainer to obtain single cell populations. These cells were then washed again in a 50 ml volume of RPMI-
35 1640 to ensure the removal of the residual DTT. The tissue was then agitated and washed as before to gather

- 70 -

the remaining iIELs. The iIELs were separated from the adipose cells and most epithelial cells on a 3-step Percol gradient, with the iIELs banding at the 40% to 80% interface. These cells were then washed twice with
5 RPMI-1640 to remove traces of Percol, immunostained with CD103 (integrin alpha IEL) antibodies, and separated on a FACs Star cell sorter. These sorted cells were then either used to prepare total RNA directly using Trizol (Gibco BRL, Gaithersburg, MD), or
10 activated overnight on plate-bound activating antibodies, which crosslink the gamma/delta TCR, alpha/beta TCR, or CD3. The RNA was prepared as above and pooled for use in constructing EST cDNA libraries.

A cDNA clone, designated smil2-00082-a1,
15 contained nucleotide sequence homology to CD28 (Figure 1B). Translation of the sequence and subsequent comparison to known proteins in a public database revealed 19% amino acid identity with murine CD28 (Figure 1B). This low homology was significant because
20 murine CD28 shares only 26% amino acid identity with murine CTLA-4. All of the putative cysteines thought to be critical for intra- and inter-molecular cysteine bonding in the CD28/CTLA-4 family were found to be conserved (amino acid residues 83, 109, and 137;
25 relative to the initiating methionine). In addition, the overall length of the putative open reading frame, and the relative position of the transmembrane domain, were similar to those of both CD28 and CTLA-4. We named the gene CRP1, for CD28-Related Protein-1.

30

Example 2

Cloning of Human CRP1 cDNA

The nucleic acid sequence encoding human CRP1
35 protein is identified by the following procedures. A human cDNA library was prepared from enriched

- 71 -

lymphocytes from peripheral human blood from normal human volunteers. The lymphocytes were purified and red blood cells were removed by Lymphocyte Separation Media (ICN Pharmaceuticals, Inc., Costa Mesa, CA). The cells were then activated overnight in media containing 10 ng/ml PMA, 500 ng/ml ionomycin, and plate-bound activating antibodies to CD3. Total RNA was prepared from the activated cells by the Trizol method (Gibco/BRL) and poly A RNA was isolated by Dynal bead purification. cDNA was made from the isolated poly A RNA and size selected for largest cDNA fragments. The size selected cDNA was then ligated into the plasmid pSPORT (Gibco/BRL). DNA encoding human CRP1 protein is obtained by screening the activated lymphocyte cDNA library by either recombinant bacteriophage plaque, or transformed bacteria colony hybridization protocols (Sambrook et al. *Supra*). The phage or plasmid cDNA library are screened using radioactively-labeled probes derived from the murine CRP1 gene clone as described in Example 1 and Figure 1. The probes are used to screen nylon filters lifted from the plated library. These filters are prehybridized for 4 hr at 42°C in 50% formamide, 5X SSPE, 2X Denhardt's solution, 0.5% SDS, and 100 µg/ml salmon sperm DNA and then hybridized for 24 hr at 42°C in 50% formamide, 5X SSPE, 2X Denhardt's solution, 0.5% SDS, 100 µg/ml salmon sperm DNA, and 5 ng/ml mB7RP1 probe. The blots are washed in 2X SSC, 0.1% SDS for 10 min at RT, 1X SSC, 0.1% SDS for 10 min at 50°C, 0.2X SSC, 0.1% SDS for 10 min at 50°C, then 0.2X SSC for 10 min at 50°C again. Inserts obtained from any human CRP1 clones are sequenced and analyzed as described in Example 1.

Example 3

B7RP1 DNA and Amino Acid Sequence

- 72 -

A cDNA clone, designated sm11-00003-g5, contained nucleotide sequence homology to B7.1 (CD80) and B7.2 (CD86). Translation of the sequence (Figure 2A) and subsequent comparison to known proteins in a public database revealed 20% amino acid identity with murine B7.1 (Figure 2B). This low homology was significant because murine B7.1 shares only 24% amino acid identity with murine B7.2. Despite this low homology, critical cysteine residues are conserved between the open reading frame of this clone and murine B7.1 and B7.2 at residues 62, 138, 185, and 242 (relative to the initiating methionine, Figure 2B). The approximate mature protein length and the location of the transmembrane region relative to the carboxy terminus are also similar in the putative ORF of this clone, as compared to B7.1 and B7.2. We named the gene B7RP1, for B7-Related Protein-1.

20

Example 4

Cloning of Human B7RP1 cDNA

A Genbank blast homology search (GCG, University of Wisconsin) using murine B7RP1 sequence (see Figure 2) retrieved a clone (AB014553) containing a 4358 bp sequence with 1679 bp of ORF. PCR cloning primers were designed according to this sequence. A DNA fragment of 1313 bp was obtained by 5' and 3' RACE using Human Lymph Node Marathon-Ready™ cDNA (Clontech, Palo Alto, CA) according to the manufacturer's recommended procedures.

30

Primers used for full length human B7RP1:

2083-75 ACC ATG CGG CTG GGC AGT CCT GGA

35

(SEQ ID NO: 25)

2083-76 TGG TGA CCT ACC ACA TCC CAC AG

- 73 -

(SEQ ID NO: 26)

2083-77 TCC GAT GTC ATT TCC TGT CTG GC

(SEQ ID NO: 27)

2083-78 GCT CTG TCT CCG GAC TCA CAG CCC

5 (SEQ ID NO: 28)

2113-29 GTG GCA GCA AAC TTC AGC GTG CCC GTC G

(SEQ ID NO: 29)

2113-30 CCC AAC GTG TAC TGG ATC AAT AAG ACG G

(SEQ ID NO: 30)

10 2113-31 GCG TGC TGA GGA TCG CAC GGA CCC CCA G

(SEQ ID NO: 31)

Primers 2083-75 and 2083-76 were used to amplify the 5' end of the gene using RACE protocols. Primers 2083-77, 2083-78, 2113-29, 2113-30, and 2113-31, were used to amplify the 3' end of the gene using RACE protocols.

The resulting nucleotide sequence contained an ORF of 288 amino acid residues beginning at the methionine. The predicted mature human B7RP1 amino acid sequence was then compared to the mature mouse B7RP1 amino acid sequence (Figure 3B) and found to share 48% amino acid identity. This homology is significant because the homology between species is low with the CD80 (B7.1) gene, in fact, the mouse and human CD80 share only 41% amino acid identity. Importantly, the human B7RP1 protein conserve critical cysteine residues necessary for Ig loop structures (amino acid residues 16, 92, 138, 194, and 195, relative to the mature protein, Figure 3B). In addition, the overall length and position of the transmembrane domain are consistent with a human B7RP1 homolog.

Example 5

Expression of B7RP1 RNA

- 74 -

RNA in situ hybridization using RNA probes to the B7RP1 gene. Adult mouse tissues were fixed in 4% paraformaldehyde, embedded in paraffin, and sectioned at 5 μ m. Prior to in situ hybridization, tissues were
5 permeabilized with 0.2M HCL, followed by digestion with Proteinase K, and acetylation with triethanolamine and acetic anhydride. Sections were hybridized overnight at 55°C with a 969 base 33 P-labeled riboprobe corresponding to nucleotides 1 to 969 of the mouse
10 B7RP1 sequence. Excess probe was removed by RNase digestion, followed by a series of washes in buffer with decreasing salt concentrations, and then a high stringency wash in 0.1X SSC at 55°C. Slides were dipped in Kodak NTB2 emulsion, exposed at 4°C for 2-3
15 weeks, developed, and counterstained with hematoxylin and eosin. Sections were examined with darkfield and transmitted light illumination to allow simultaneous evaluation of the tissue morphology and the hybridization signal.

20 The analysis of the B7RP1 RNA by in situ hybridization showed that the B7RP1 RNA was highly expressed in areas of lymphoid maturation and lymphocyte activation. B7RP1 RNA was expressed in the lymphoid tissues of the thymus, Peyer's patches of the
25 intestine, spleen, and lymph nodes. Expression within these lymphoid tissues demonstrated that the B7RP1 RNA was generally expressed in the areas of B-cell and other APC involvement. These regions include the medulla area of the thymus, the primary follicles of
30 the lymph nodes, and the follicular and dome regions of the Peyer's patches. The expression of B7RP1 RNA is highly specific to the regions of APC involvement in lymphoid tissues.

35 The analysis of several non-lymphoid tissues also revealed B7RP1 expression in regions of APC

- 75 -

involvement. In the lung, B7RP1 expression was found in the submucosal regions, consistent with a function in antigen processing. In the small intestine, B7RP1 RNA was found in the lamina propria. Notably, we found
5 a section of damaged liver, which showed lymphocyte infiltration that overlapped with the expression of B7RP1 RNA. This coincidence of B7RP1 expression with lymphocyte accumulation in response to tissue damage strongly indicates that B7RP1 is involved in lymphocyte
10 activation.

Example 6

Expression of CRP1 RNA

15 RNA in situ hybridization using RNA probes to the CRP1 gene. Mouse tissues were prepared as in Example 5. Tissue permeabilization, probe hybridization, slide treatment, and tissue staining were as described in Example 5. Sections were
20 hybridized overnight at 55°C with a 603 base ³³P-labeled riboprobe corresponding to nucleotides 1 to 603 of the mouse CRP1 sequence. Sections were examined with darkfield and transmitted light illumination to allow simultaneous evaluation of the tissue morphology
25 and the hybridization signal.

Lymph nodes from normal mice or a mouse treated with oxazolone were sectioned and analyzed for CRP1 RNA expression. The sensitized mouse lymph node showed greater expression of CRP1 RNA than the normal
30 mouse lymph node. The expression of CRP1 was in the paracortex, a region of T-cell activity. Therefore, the expression of CRP1 RNA is consistent with that of T-lymphocyte expression and is up-regulated upon T-cell activation.

- 76 -

Example 7Expression and Purification of
CRP1-Fc and B7RP1-Fc Fusion Proteins

5

To construct the DNA expression vector for the CRP1- Fc fusion protein, the coding sequence for the first amino terminal 147 amino acids of the CRP1 was fused, inframe, to the coding sequence for the carboxy terminal 235 amino acids of the human Fc gene (isotype IgG1) and ligated within the polylinker sequence of pcDNA3 (pcDNA3/CRP1-Fc). To construct the DNA expression vector for the B7RP1- Fc fusion protein, the coding sequence for the first amino terminal 269 amino acids of the B7RP1 was fused, inframe, to the coding sequence for the carboxy terminal 235 amino acids of the human Fc gene (isotype IgG1) and ligated within the polylinker sequence of pcDNA3 (pcDNA3/B7RP1-Fc). The coding sequences of both CRP1 and B7RP1 contained sequences from the N-terminus of each protein up to, but not including, the putative transmembrane region of each protein. 293T cells were transfected with either pcDNA3/CRP1-Fc or pcDNA3/B7RP1-Fc using the FuGene 6 transfection reagent (Roche Molecular Biochemicals, Indianapolis, IN). After four days, the conditioned media were collected and the Fc fusion proteins were purified by batch chromatography using Protein A Sepharose (Pharmacia). Fc fusion proteins bound to the column were eluted with three column volumes of Immunopure Gentle Elution Buffer (Pierce), and then were dialyzed against 150 volumes of 20 mM HEPES, 100 mM NaCl, pH 7.5. The dialyzed protein was concentrated using Macrosep centrifugal concentrates, 30 kD MWCO (Pall Filtron), and the protein concentrations were calculated using extinction coefficients derived from the amino acid sequence of

- 77 -

each protein. Expression of CRP1-Fc fusion protein is shown in Figure 4A, expression of B7CPR1-Fc fusion protein is shown in Figure 4B.

5

Example 8

Identification of CRP1 and B7RP1
as a Receptor-ligand Pair

In order to determine whether the novel
10 proteins were part of the same costimulatory pathway as that containing CD28, CTLA-4, B7.1, and B7.2, we utilized a cell surface display assay. This assay uses ACAS (Adherent Cell Analysis and Sorting) analysis to analyze whether membrane-bound proteins expressed in
15 cells interact with various Fc fusion proteins. Cells expressing membrane-bound proteins, indicated on the left side of Figure 5, were incubated with Fc fusions proteins, indicated at the top of the figure.

Cos-7 cells, grown in DMEM media with 10%
20 FBS, were plated at 500,000 cells/well in a 24-well plate. Cells were transfected using the FuGene 6 reagent (Roche Molecular Biochemicals, Indianapolis, IN). For each transfection, 3 µl of FuGene 6 reagent was added to 47 µl of serum free DMEM media. After a
25 10 min incubation at room temperature, the mix was added to 0.25 µg of plasmid dropwise and then was incubated for 15 minutes. The above mix was then added to the cells with 0.5 ml of DMEM with 10% FBS. The cells were incubated at 37°C in a 5% CO2 atmosphere.
30 As a control, CHO D- cells, stably transfected with an expression plasmid containing the cDNA for human CD28, were also plated at 500,000 cells/well in a 24-well plate.

After 48 hr, the medium with transfection
35 reagent was removed and the cells were washed twice

- 78 -

with RPMI plus 5% FBS. 10 to 20 ng of purified Fc fusion proteins in 1 ml of media were added to the cells, which were incubated for 30 min on ice. The cells were washed three times with RPMI plus 5% FBS and then were incubated with 2 μ l of FITC-conjugated anti-human Fc antibody (1 mg/ml) for another 30 min on ice. After three successive washes with RPMI, the cells were covered with 250 μ l of RPMI media without phenol red for ACAS analysis.

ACAS analysis of the cells that bound the various Fc fusion proteins demonstrated that the B7RP1 protein bound CRP1, but not the proteins in the known costimulatory pathway, CD28 or CTLA-4. Conversely, CRP1 interacted with B7RP1, but not B7.2, a component in the known pathway. (See Figure 5). These results strongly indicate that CRP1 and B7RP1 represent a novel receptor-ligand pair, analogous to CD28 and B7.2. However, since CRP1 and B7RP1 do not interact with B7.2, CTLA-4, or CD28, they are separate and independent of the known costimulatory pathway.

Example 9

Identification of Cells Expressing B7RP1 Receptors

The B7RP1-Fc fusion protein was utilized to detect cells that expressed receptors to B7RP1, presumably including the CRP1 protein (see Example 6), by FACS analysis. Spleens were removed from female C57/Black 6 mice, ground on 100 micron mesh filters to release the lymphocytes, passed through 70 micron filters, and then washed in 50 ml of RPMI-1640. They were pelleted at 1500 rpm, resuspended in fresh RPMI, mixed to break up the clumping cells, and passed through a 40 micron filter. T-cells to be activated were seeded into 6 well plates in RPMI-1640, 5% FBS,

- 79 -

1XPSG, PMA, ionomycin, and incubated at 37°C, 5% CO₂ overnight. T-cell activation was checked by visual confirmation after 12 hr.

Activated spleen cells for immunostaining
5 were washed in PBS, 0.5% BSA (Path-o-cyte 4, ICN Pharmaceuticals) wash buffer, resuspended, and then aliquoted in 100 µl volumes. 15 µg/ml of either the CRP1-Fc fusion protein or the B7RP1-Fc fusion protein was added (1.5 µg/sample) as appropriate, and then the
10 mixtures were incubated on ice for 30 min with occasional mixing. The cells were washed twice in 5.0 ml of wash buffer. Binding of the fusion proteins was visualized with 2 µg of goat-anti-human (GaHuFc-FITC) conjugated secondary antibody in a 100 µl volume for
15 cell staining. Cell marker antibodies conjugated with PE were added with the GaHUFc-FITC, as well as control isotype-PE conjugated antibody controls where indicated (rat isotype). The samples were incubated on ice and washed as before. Visualization was done by FACScan
20 analysis with gating on the lymphocyte populations. Double staining with CD4+ antibodies and the B7RP1-Fc fusion protein indicated that the cells expressed both the CD4 marker and the receptor to B7RP1, presumably CRP1 (Figure 6). Similarly, double staining with CD8+
25 antibodies and the B7RP1-Fc fusion protein demonstrated that cells expressed both CD8 and B7RP1 receptors (Figure 6). We could not reliably detect such double staining cells in inactivated splenocyte preparations. Since CD4 and CD8 are T-lymphocyte markers, we can
30 postulate that CRP1 is expressed on activated CD4+ and CD8+ T-cells. These data are consistent with the increased expression of CRP1 RNA in the T-cell regions of lymph nodes from sensitized mice as compared to normal mice (Example 6).

35

Example 10

- 80 -

Identification of Cells Expressing CRP1 Ligands

The CRP1-Fc fusion protein was utilized to detect cells that expressed ligands to CRP1, presumably including the B7RP1 protein (see Example 8), by FACS analysis (Figure 7). Splenocytes were prepared as in Example 8, except the 12 hr T-cell activation step was omitted and the cells were directly analyzed. Splenocytes were double stained with CD45R (B220) marker antibodies and the CRP1-Fc fusion protein. Cells were detected that expressed both the CD45R B-cell marker and the putative ligands for CRP1, presumably including B7RP1 (Example 8). Therefore, we conclude that B7RP1 is expressed on B-cells, a type of antigen-presenting cell. These data are consistent with the expression of B7RP1 RNA in B-cell regions of various lymphoid tissues (Example 5).

FACS analysis of the expression of B7RP1 on peritoneal macrophages (Figure 8). Peritoneal cells were collected by local lavage from a normal mouse and washed before being incubated with the CRP1-Fc fusion protein or the Fc protein as a control or with the F4/80 monoclonal antibody (which detects an antigen specific for macrophages) or an irrelevant, isotype-matched control monoclonal antibody. Cells were then washed again and incubated with goat-anti-human Fc-FITC conjugated antibody. After further washing, cells were assessed in a FACS analyzer for their light scattering and fluorescence staining properties. Peritoneal cells were first distinguished in subsets on the ground of their light scattering properties (Figure 8A). Macrophages were identified in region 5 (R5) because of their ability to strongly scatter light forward (FSC) and sideways (SSC) and because of their positive staining for the F4/80 antigen, a marker for macrophages (Figure 8B). Macrophages in region 6 (R6)

- 81 -

were singled out on the basis of their less intense staining for the F4/80 antigen and found to be stained by the CRP1-Fc fusion protein (Figure 8C). These data indicate that ligands for CRP1, possibly including
5 B7RP1, are expressed on macrophages, a professional antigen presenting cell. This is consistent with CRP1 and B7RP1 function in T-lymphocyte activation.

Example 11

10 In vitro Inhibitory Activity of the B7RP1-Fc Fusion Protein on ConA-Stimulated T-Lymphocytes

Mouse splenocytes were prepared as in Example 8 and enriched for T-lymphocytes by negative selection
15 (R and D Systems, Inc., Minneapolis, MN)). 200,000 splenocytes were used in T-cell proliferation assays in a 96-well round-bottom plate. Cells were incubated for 1 hr with media (no adds), CRP1-Fc, B7RP1-Fc, or B7.2-Fc, fusion proteins as indicated Figure 9. Media (no
20 adds), or Con A at various concentrations were added as indicated in at the bottom of Figure 9. The cells were then incubated at 37°C and 5% CO₂. After 42 hr, cells were pulsed with 3H-thymidine for 6 hr, harvested and incorporated radioactivity determined. Average CPM and
25 standard deviation from triplicate samples are represented in Figure 9.

The Fc fusion proteins did not demonstrate significant T-cell stimulatory or inhibitory activity by themselves, however, in the presence of 1 µg/ml and
30 3 µg/ml Con A, both the B7RP1-Fc and the known B7.2-Fc fusion proteins showed significant inhibitory activity (Figure 9). At high concentrations (10 µg/ml), Con A stimulation results in cell death, presumably through over-activation of the T-cells. Addition of either
35 B7RP1-Fc or B7.2-Fc, significantly protected the cells from the detrimental effects of high concentrations of

- 82 -

Con A. In both inhibitory and protective functions, the effect by B7RP1-Fc protein was greater than B7.2-Fc protein on the Con A stimulated cells. These data indicate that the B7RP1 protein functions to regulate
5 T-cell proliferation.

Example 12

Systemic delivery of B7RP1-Fc fusion protein in transgenic mice

10

The B7RP1-Fc fusion protein described in Example 7 was subcloned into an ApoE-liver specific expression vector (Simonet et al. J. Clin. Invest. 94, 1310-1319 (1994) and PCT Application No. US94/11675).
15 The coding region was excised from pCEP4/B7RP1-Fc using the restriction enzymes, Spe I and Not I, and the fragment subcloned into the same sites in the previously mentioned ApoE-liver specific expression vector. The resultant plasmid, HE-B7RP1-Fc, was
20 sequenced through it's protein coding region, and sequences flanking the coding region, to ensure it was mutation free.

The plasmid was amplified and purified through two rounds of CsCl density gradient
25 centrifugation. The purified plasmid DNA was digested with the restriction enzymes, Cla I and Ase I, and the 1.5 kb transgene insert was purified by agarose gel electrophoresis. The purified fragment was diluted to a stock injection solution of 1 µg/ml in 5 mM Tris, pH
30 7.4, and 0.2 mM EDTA. Single-cell embryos from BDF1 X BDF1-bred mice were injected essentially as described (Brinster et al., Proc. Natl. Acad. Sci. USA 82, 4338 (1985)), except that injection needles were beveled and siliconized before use. Embryos were cultured
35 overnight in a CO2 incubator and 15 to 20 2-cell

- 83 -

embryos were transferred to the oviducts of pseudopregnant CD1 female mice.

Following term pregnancy, 56 offspring were obtained from implantation on the microinjected embryos. The offspring were screened by PCR amplification of the integrated transgene in genomic DNA samples. The target region for amplification was a 369 bp region of the human Apo E intron which was included in the expression vector. The oligos used for PCR amplification were:

5'-GCC TCT AGA AAG AGC TGG GAC-3' (SEQ ID NO: 32)

5'-CGC CGT GTT CCA TTT ATG AGC-3' (SEQ ID NO: 33)

The conditions for the PCR were: 94°C for 2 min, 1 cycle; 94°C for 1 min, 63°C for 20 sec, and 72°C for 30 sec, 30 cycles. Of the 56 original offspring, 7 were identified as PCR positive transgenic founder mice.

At 12 weeks of age, nine transgenic founders (mouse #1, 2, 4, 6, 8, 30, 32, 33, 40) and five controls (mouse #5, 9, 10, 25, 28) were sacrificed for necropsy and pathological analysis. Total cellular RNA was isolated from the livers of the founder animals and negative control littermates as described (McDonald et al. Meth. Enzymol. 152, 219 (1987)). Northern blot analysis was performed on these samples to assess the level of transgene expression. Approximately 10 µg of total RNA from each animal was resolved by agarose electrophoresis denaturing gels (Ogden et al. Meth. Enzymol. 152, 61 (1987)), then transferred to HYBOND-N nylon membrane (Amersham), and probed with ³²P dCTP-labeled mB7RP1-Fc insert DNA. Hybridization was performed for 1 hr at 63°C in ExpressHyb Solution (Clonetech) and 2-4 X 10⁶ CPM of labeled probe/ml

- 84 -

hybridization buffer. Following hybridization, blots were washed twice in 2X SSC, 0.1% SDS at room temperature for 5 min each, and then twice in 0.1X SSC, 0.1% SDS at 55°C for 15-20 min each. Expression of the transgene in founder and control littermates was determined following autoradiography.

Northern blot analyses indicated that seven of the transgenic founders expressed detectable levels of the transgene RNA (mouse #1, 2, 6, 8, 32, 33, and 40). The negative control mice and three founders (#4, 30, and 31) did not express detectable levels of RNA. Since the B7RP1-Fc fusion protein was determined to be secreted from mammalian cells in culture (Figure 4B and Example 7), expression of the transgene mRNA should be indicative of the level of systemically delivered gene product.

Example 13

Biological activity of B7RP1-Fc fusion protein

20

Seven of the transgenic mice (mouse #1, 2, 6, 8, 32, 33, and 40) and five control littermates (#5, 9, 10, 25, and 28) were sacrificed for necropsy and pathological analysis using the following procedures: Prior to euthanasia, all animals had their identification numbers verified, then were weighed, anesthetized and blood drawn. The blood was saved as both serum and whole blood for a complete serum chemistry and hematology panel. Radiography was performed just after terminal anesthesia by lethal CO₂ inhalation, and prior to gross dissection. Tissues were then removed and fixed in 10% buffered Zn-formalin for histological examination. The tissues collected included the liver, spleen, pancreas, stomach, duodenum, ileum, Peyer's patches, colon, kidney, reproductive organs, skin, mammary glands, bone, brain,

- 85 -

heart, lung, thymus, trachea, esophagus, thyroid/parathyroid glands, jejunum, cecum, rectum, adrenal glands, white and brown fat, sciatic nerve, bone marrow, urinary bladder, and skeletal muscle.

5 Prior to fixation, the whole organ weights were determined for the liver, heart, stomach, kidney, adrenals, spleen, and thymus. After fixation, the tissues were processed into paraffin blocks, and 3 μ m sections were obtained.

10 Immunohistochemistry for the B-lymphocyte marker, B220, and the T-lymphocyte marker, CD3, was performed. To detect B220 or CD3 expression, formalin fixed, paraffin embedded, 4 μ m sections were deparaffinized and hydrated to deionized water. The
15 sections were quenched with 3% hydrogen peroxide, blocked with Protein Block (Lipshaw, Pittsburgh, PA), and incubated in rat monoclonal antibody to B220 (Pharmingen, San Diego, CA) or rabbit polyclonal antibody to CD3 (Dako, Carpinteria, CA). The
20 antibodies were detected by biotinylated rabbit anti-rat or goat anti-rabbit immunoglobulins, peroxidase conjugated streptavidin, (BioGenex, San Ramon, CA) with DAB as chromagen (Biotek, Santa Barbara, CA). Sections were counterstained with hematoxylin.

25 In this study, normal clinical signs were reported during the in-life phase of the study. The whole body radiographs of the transgenic mice were comparable to those of the control mice. The overall hematologic parameters of the transgenic mice were
30 comparable to those of the negative control group, although sporadic changes in individual mice were present: transgenic #8 and #40 had increased serum globulin levels (hyperglobulinemia) and #32 and #33 had globulin levels in the high normal range accompanied by
35 albumin levels in the low normal range, which is a pattern commonly seen with chronic antigenic

- 86 -

stimulation of the immune system. Organ weights of the other transgenic mice were not significantly different from those of the control group.

The following histopathological changes were present in the transgenic mice: The mesenteric lymph nodes of the transgenic B7RP1-Fc mice were moderately to markedly enlarged when compared to the control mice (Fig. 10A-10D; Fig. 11A-11E). The cortex had prominent follicular hyperplasia seen as enlarged secondary follicles (Fig. 10B-11B) with large germinal centers containing mostly B220+ B cells (Fig. 11D) and a few scattered CD3+ T cells (Fig. 11F). The paracortical (CD3+ T cell) area was also moderately enlarged (Fig. 11B-11F) and the medullary sinuses had slightly increased numbers of fleshy macrophages (sinus histiocytosis). The most conspicuous change in the nodes was present in the medullary cords, which were mildly to markedly expanded by large numbers of well-differentiated plasma cells in the B7RP1-Fc transgenic mice (Fig. 10D). In transgenic mouse #40, small numbers of scattered Russell bodies (i.e. plasma cells with prominent, large, round, intracytoplasmic vesicles containing immunoglobulins) were also found in the medullary cords (Fig. 10D). Interestingly, the other internal and peripheral lymph nodes (e.g. cervical, inguinal) had similar morphologic features of reactive lymphoid hyperplasia suggestive of a systemic response. These findings are consistent with a chronic, ongoing immune stimulation with enhancement of the humoral immune reaction, which leads to B cell proliferation and terminal differentiation into plasma cells.

The spleen of B7RP1-Fc transgenic mice had variably enlarged white pulp areas with moderate reactive lymphoid hyperplasia involving particularly the B-cell secondary follicles with prominent germinal

- 87 -

centers and periarteriolar T-cell sheaths when compared to the control mice (Fig. 10E-10F). Another conspicuous finding in B7RP1-Fc transgenic mice was minimal to mild plasmacytosis in the marginal zone surrounding the white pulp areas and in the adjacent red pulp. Transgenic mouse #6 had a few scattered Russell bodies (Fig. 10F, inset). The red pulp had mild to moderate extramedullary hematopoiesis, which was comparable to that seen in the control mice (Fig. 10E).

The small intestinal Peyer's patches were mildly to markedly enlarged in the B7RP1-Fc transgenic mice over those of the control mice (Fig. 10G) and had very large follicles with prominent germinal centers, particularly in transgenic mouse #40 and #32 (Fig. 10H). In addition, there was a minimal (in #32) to mild (in #8 and #33) increase in the numbers of lymphocytes and plasma cells (admixed with a mild eosinophil infiltrate in the ileum of mouse #32) in the thickened lamina propria layer of the mucosa, which was present in the small intestine, but more prominent in the colon of the transgenic mice. The large intestinal lymphoid aggregates (GALT) were also slightly more prominent in some B7RP1-Fc transgenic mice (particularly mouse #8 and #2) than in the control group.

Generally, the other tissues examined, including the thymus, bone marrow, liver, lung, heart, pancreas, kidneys, adrenal gland, thyroid, parathyroid, trachea, reproductive organs, urinary bladder, mammary gland, skin, skeletal muscle, peripheral nerve, brain, esophagus, stomach, small and large intestine, bone (femur/tibia), stifle joint, white and brown fat appeared normal and comparable to the background changes detected in the control mice.

- 88 -

The data from this study demonstrate that overexpression of the B7-related protein Fc chimera (B7RP1-Fc) in transgenic mice induces a phenotype characterized by prominent reactive lymphoid hyperplasia detected in the spleen, peripheral and internal lymph nodes, and gut-associated lymphoid tissue, as follicular hyperplasia, expansion of T cell areas and conspicuous plasmacytosis accompanied by hyperglobulinemia in some animals. The plasmacytosis is accompanied by higher levels of circulating IgG (mean \pm SD = 597 \pm 298 mg/ml in transgenic mice vs. 209 \pm 80 mg/ml in control littermates, n = 7, P < 0.05, t test), in particular IgG2a (217 \pm 100 mg/ml vs. 75 \pm 29 mg/ml, n = 7, P < 0.01, t test). The induction of IgG2a is normally associated with a Th1 cytokines such as IFN- γ . Thus, B7RP-1 induces B- and T-cell proliferation and stimulates B-cells to differentiate into plasma cells and to produce immunoglobulin.

These changes are consistent with a persistent systemic immune response with hyperstimulation of the humoral arm of the immune system which results in B cell stimulation, proliferation, and differentiation to antibody-producing plasma cells throughout the lymphoid organs examined.

We conclude from the marked lymphoid hyperplasia demonstrated in the B7RP1-Fc transgenic mice that B7RP1 protein has significant *in vivo* biological activity, related to immune system stimulation.

Example 14

Cloning of human B7RP1

Normal human circulating peripheral lymphocytes were separated from red blood cells using

- 89 -

Lymphocyte Separation Medium (ICN Pharmaceuticals). The T-cells were then activated with 10 µg/ml plate bound anti-CD3 antibody (Immunotech, Westbrook, ME), 10 ng/ml PMA, and 500 ng/ml ionomycin overnight (16 hours) at 5 37°C and 5% CO₂. Total RNA was then prepared from the cells using TRIzol reagent (Gibco BRL). The cells were pelleted by centrifugation and the cell pellet was resuspended in 1 ml TRIzol reagent for each 5 X 10⁶ cells and incubated at room temperature for 5 min. 0.2 10 ml chloroform per 1 ml original TRIzol reagent was then added. The tubes were shaken vigorously by hand for 15 seconds and incubated for 3 minutes at RT and centrifuged at 13,000rpm for 15 min at 4°C. Following centrifugation, the clear upper aqueous phase which 15 contains the RNA was collected and the sample RNA was precipitated by the addition of isopropyl alcohol. The solution was then incubated at RT for 10 min, the RNA pelleted, washed with 75% ethanol, and then centrifuged at 15,000 rpm for 5 min at 4°C. The pellet was air 20 dried, resuspend in RNase-free water, then aliquoted, and stored at -80°C until later use.

The library was constructed using the SuperScript Plasmid System for cDNA Synthesis and Plasmid Cloning (Gibco BRL). Briefly, cDNA inserts 25 with an average size of 2 kb, were ligated into the pSport vector at SalI/ NotI cloning site. The ligated plasmids were electroporated into Electromax transformation competent *E.coli* (Gibco BRL), titered and plated at fifteen thousand colonies per LB plate 30 (ampicillin 100 µg/ml). 300,000 colonies were lifted onto colony/plaque screen hybridization transfer membranes (NEN Life Sciences), denatured in 0.5 N NaOH, 1.5 M NaCl for 5 minutes, then neutralized successively for 5 minutes each in the following buffers, 1 M Tris 35 HCl pH 8.0, 0.5 M Tris HCl pH 8.0 and 1.5 M NaCl and 2X SSC. The filters were then crosslinked by

- 90 -

ultraviolet irradiation and baked for 30 min at 80°C in a vacuum oven. The filters were pre-washed extensively in 2X SSC at 42°C to remove debris, then prehybridized at 42°C in 50% formamide, 5X SSPE, 5X Denhardt's solution, 0.5% SDS, 100 µg/ml salmon sperm DNA, for 2 hours.

The human lymphocyte cDNA library was screened with an 895 bp DNA fragment having nucleotides 1-711 as shown in Figure 3A, 167 bps immediately 5' to the initiator methionine codon in Figure 3A and 17 bps immediately 3' to position 711 in Figure 3A. This upstream 5' sequence of 167 base pairs was obtained by 5' RACE of the HuB7RP1 cDNA (Example 4) and was released from a TOPO TA vector (Invitrogen, Carlsbad, CA) at the Eco RI restriction enzyme cleavage site. This insert was twice purified on a 0.8% agarose TAE gel. A DNA gel purification kit (Qiagen) was used to isolate the DNA insert from the agarose.

125 ng of the DNA fragment was labeled with ³²P dCTP (Amersham) following the Redi-Prime 2 (Amersham) random prime labeling system protocol. The colony lift filters were then allowed to hybridize with the probe at 42°C in the following buffer overnight at 42°C; 50% formamide, 5X SSPE, 2X Denhardt's solution, 0.5% SDS, 100 mg/ml ssDNA. The specific activity of the probe was 2.38 X 10⁹ cpm/µg DNA, in approximately 2 ng labeled probe per ml hybridization buffer. The probe was removed and saved for the next round of screening. The filters were then washed in 2X SSC, 0.1% SDS RT for 15 min, followed by 1X SSC, 0.1% SDS at 55°C for 15 min, and 1X SSC, 0.1% SDS at 60°C for 10 min. The filters were wrapped in plastic and exposed to autoradiography film overnight at -80°C with 2 enhancing screens. Three independent positive clones were identified. Exposures were aligned to the

- 91 -

bacterial plates and the positive clones scraped, diluted and replated on LB plates with ampicillin 100µg/ml, grown overnight as before and the colonies were lifted, prepared, and probed as described previously. Three independent clone colonies were isolated, the DNA was isolated, and DNA sequenced for each clone in triplicate.

The full-length of the human B7RP1 protein is 302 amino acids. The polypeptide length and relative position of the transmembrane domain, is consistent with other B7 family members. The human B7RP1 gene has 43% amino acid identity with the mouse clone. This degree of homology is significant since the mouse and human CD80 proteins are only 41% identical. Notably conserved between the mouse and human genes are the cysteine residues at amino acid positions 37, 113, 158, 215, and 216.

Example 15

Cloning of human CRP1

A Genbank blast homology search (GCG, University of Wisconsin) using murine B7RP1 sequence (see Figure 2) retrieved a genomic clone (Gen Bank Assession NO. AQ022676) containing a 104 bp sequence that showed high homology with the murine CRP1 gene. PCR cloning primers were designed to overlap this sequence.

5'-GCA TAT TTA TGA ATC CCA-3' (SEQ ID NO: 34)
5'-ACT ATT AGG GTC ATG CAC-3' (SEQ ID NO: 35)

Using the above primers, a 151 bp DNA fragment of the murine CRP1 was PCR amplified using the murine CRP1 plasmid described in Figure 1 and Example 1 as template. 125 ng of the DNA was labeled with 32P

- 92 -

dCTP (Amersham) following the Redi-Prime 2 (Amersham) random prime labeling system protocol. The colony lift filters from human peripheral blood libraries described in Example 15 were then allowed to hybridize with the probe in the following hybridization buffer overnight (15 hr) at 41°C, 50% formamide, 5X SSPE, 2X Denhardt's solution, 0.5% SDS, 100 µg/ml ssDNA. The specific activity of the probe was 3.52×10^9 cpm/µg DNA, 1.5 ng labeled probe/ml hybridization buffer. The probe was pulled off and saved for the next round of screening. The filters were then washed in 2X SSC, 0.1% SDS at RT for 10 min, followed by 1X SSC, 0.1% SDS at 37°C for 7 minutes, 40°C for 7 minutes, 44°C for 7 minutes, then 50°C for 7 minutes, continually monitoring the rate at which the filters were releasing the labeled probe. The filters were wrapped in plastic and exposed to film overnight at -80°C with 2 enhancing screens. This method revealed 9 possible independent positive clones. Exposures were aligned to the bacterial plates and the positive clones scraped, deposited into 200 µl SOC, 2 serial dilutions of 1:10 were performed and 70 µl from the second dilution was replated on LB plates containing ampicillin at 100 µg/ml and grown overnight as before. The colonies were lifted, prepared and probed as before. Eight independent clones were isolated and DNA prepared by the Qiagen miniprep method.

A cDNA clone containing an open reading frame of 199 amino acids was obtained (Figure 13A). This cDNA clone contained nucleotide and amino acid homologies to the murine CRP1 clone described in Example 1 and Figure 1. The nucleotides corresponding to the open reading frame of this human clone was 77% identical to the murine CRP1 gene. Translation of the human sequence and subsequent comparison with the

- 93 -

murine CRP1 protein revealed 69% amino acid identity with the murine protein (Figure 13B). In addition, the motif between amino acids 114 to 119, "FDPPPF", was conserved between the murine and human CRP1 genes.

5 This motif corresponds to the "MYPPPY" motif in murine and human CD28 that is essential for B7 protein interaction. Furthermore, the cysteines at amino acid positions 42, 109, and 141 are also conserved. These cysteines correspond to cysteines in CD28 and CTLA-4 at
10 are involved in Ig loop formation and intermolecular disulfide dimerization. The close similarity with murine CRP1, and structural similarities with the CD28 homology family, indicate that this is the human CRP1 homolog.

15

Example 16

CRP-1 is expressed on resting memory T-lymphocytes

20 In order to study CRP-1 expression on memory T-cells, splenic T-cells were collected from 6-7 month old mice. These cells were double-stained using B7RP-1-Fc labeled by an FITC-conjugated anti-human Fc antibody and a PE-conjugated antibody to either CD44,
25 CD45RB, or CD69. Staining with the B7RP-1-Fc fusion protein detects expression of CRP-1 protein on these T-cells. Older mice show more CRP-1+ splenic T-cells than younger mice. Interestingly, a conspicuous number of these cells are CD44 high(Fig 14a) and CD45RB low
30 (Fig 14b), a profile typical of memory T-cells. These CRP-1+ memory T-cells are in a resting state, since they do not express the activation marker CD69 (Fig. 14c). The expression of CRP-1 on memory T-cells indicates that CRP-1 has costimulatory functions on
35 memory T-cells.

- 94 -

Example 17

In vitro T-cell costimulation
inhibited by antibodies to B7RP-1.

5 To determine if the B7RP-1 protein has
functional relevance to T-cells, we incubated CD3+ T-
cells with the B7RP-1-Fc fusion protein and an anti-CD3
antibody in an *in vitro* proliferation assay. Rabbit
anti-mouse B7RP1 polyclonal antibodies or rat anti-
10 mouse B7RP1 monoclonal antibodies were then used to
specifically inhibit B7RP1-Fc costimulated
proliferation *in vitro*.

B7RP-1 rabbit polyclonal antiserum preparation

15 Three New Zealand white rabbits (5-8 lbs.
initial weight) were injected IM with murine B7RP1
protein. Each rabbit was immunized on day 1 with 150 µg
of murine B7RP1 protein emulsified in an equal volume
of Hunters Titer Max complete adjuvant. Further boosts
20 (days 14 and 28) were performed by the same procedure.
Antibody titers were monitored by EIA. After the
second boost, the antisera revealed moderate antibody
titers. A 30ml production bleed was then obtained from
each animal. This was repeated each week for 6 weeks.
25 Polyclonal antibodies were then purified by protein-A
agarose chromatography, followed by negative selection
Fc protein affinity chromatography and positive
selection by B7RP-1-Fc affinity chromatography.

30 Rat anti-murine B7RP1 monoclonal antibody preparation

Rat anti-murine B7RP1 monoclonal antibodies
were generated as described in Practical Immunology,
second edition (1980; L. Hudson and F.C. Hay; Blackwell
Scientific Publications; St. Louis, MO). Briefly, Lou
35 rats (Harlan; Indianapolis, IN) were injected

- 95 -

intraperitoneally with muB7RP1-Fc fusion protein emulsified in Freund's Adjuvant at 4 week intervals. Three days prior to fusion, rats were boosted intravenously with soluble muB7RP1. On the day of fusion, the animal was sacrificed under carbon dioxide and the spleen removed aseptically. Single cell suspension was generated using a tissue stomacher. Both splenocytes and Y3-Ag1.2.3 myeloma cells (American Type Culture Collection; Rockville, MD) were washed in serum-free media then fused by the addition of polyethylene glycol (PEG 1500; Boehringer Mannheim Biochemicals; Indianapolis, IN). The cells were rinsed once, resuspended in serum-containing media, and plated into 96-well tissue culture plates. Ten to 12 days later, media from each well was tested for specific antibody to B7RP1 via a direct Enzyme-linked Immunosorbent Assay (EIA). Cells from wells indicating potential binding were grown to 10 ml cultures and frozen in liquid nitrogen. Media from each culture was further tested in flow cytometry and in a functional T-cell proliferation assay. Those determined to be of interest by these methods were plated to single cell colonies, selected again by EIA, and final cell lines maintained for antibody generation. Antibodies were purified from the cell media by protein A agarose chromatography.

T-cell preparation and T-cell proliferation assay

T-cells from the spleens of C57Bl/6 mice (8-12 week-old females, Charles River Laboratories) were purified by negative selection through a murine T-cell enrichment column (R&D Systems). The T-cells were then either used directly or further purified by antibody and complement lysis as follows. Cells were resuspended (2.5×10^6 cells/ml) in RPMI medium containing antibodies (all at 10 μ g/ml and from

- 96 -

Pharmingen) against murine CD11b (Clone M1/70), NK-1.1 (Clone PK136), CD8a (Clone 53-6.7), I-A^b (Clone M5/114.15.2), CD11c (Clone HL3), and the B220 antigen (Clone RA3-6B2). The cells were then incubated on ice for 30 min, pelleted at 1200 rpm, resuspended in 4:1 vol/vol of RPMI: rabbit complement (Sigma, #S-7764), and incubated for an additional 30 min at 37°C. The cells were pelleted again, and the complement treatment was repeated. Before plating, the cells were washed with RPMI containing 10% FCS. U-bottomed 96 well plates were coated with an anti-CD3 antibody (Clone 145-2C11, Pharmingen) at concentrations ranging between 0 and 1.2 µg/ml, and anti-human IgG Fab₂ (Sigma, 12.5 µg/ml) overnight at 4°C, followed by a 6-9 hr incubation at 37°C. T-cells (1x 10⁵/well) were cultured in the absence or presence of various Fc fusion proteins for 48 hr and were pulsed during the last 18 hours with 1 µCi of ³H-thymidine. Control Fc proteins included a fusion protein of OPG and Fc and a nonfused Fc protein fragment. The cells were then harvested and the incorporated radioactivity was counted. B7RP-1-Fc co-stimulates T-cells to proliferate in a dose-dependent fashion (Fig. 15a), and an anti-B7RP-1-Fc antibody specifically inhibits this co-stimulation dose-dependently (Fig. 15b).

Example 18

Inhibitors of the CRP-1/B7RP-1 pathway decrease the onset of rheumatoid arthritis induced by collagen.

Collagen-induced arthritis (CIA) is an animal model of autoimmune polyarthritis in rodents and primates that has many similarities with rheumatoid arthritis in humans. Immunization with heterologous

- 97 -

species of type II collagen (CII) induces an autoimmune response to CII that leads to the development of CIA in susceptible mouse strains. Congenic strains of mice with H-2^r and H-2^a are highly susceptible to CIA. CIA is mediated by the synergistic effects of both CII-reactive T cells and antibodies. Porcine CII (Nabozny et al., Autoimmunity 20, 51-58 (1995)) was dissolved in 0.01N acetic acid at a concentration of 2mg/ml and then was emulsified at a 1:1 ratio with CFA (Difco).

Arthritis susceptible B10.RIII (H-2^r) mice (Jackson Laboratories, Bar Harbor, ME) were immunized with 100 µl of emulsion intradermally at the base of the tail. Mice were monitored 2-3 times per week for the development of arthritis. Arthritis severity was determined using a grading system for each paw as follows: 0: no arthritis; 1: redness or swelling in 1-3 toes; 2: severe swelling of paw; 3: joint ankylosis. The score of each limb was summed to give a severity range from 0 to 12 for each animal.

Mice were injected with 100ug (in 200 µL) of protein intraperitoneally twice per week. The treatment was begun 1 day after immunization with porcine CII and was stopped at day 52 post-immunization. The experiment was conducted in treatment groups of 10 mice, and animals with scores of 1 or above were scored as positive. The results are shown in Figure 16 and Table 1.

TABLE 1

Effect of CRP-1, B7RP-1, CTLA-4, and B7.2 Fc fusion proteins on the onset of arthritis

Treatment groupsMean+/-s.d. day of onset

CTLA4-Fc

60.0±0.0

- 98 -

	CRP1-Fc	48.9±13.2
	B7.2-Fc	28.4±14.1
	B7RP1-Fc	33.9±16.6
5	PBS	37.7±17.1

10 In mice treated with CRP1-Fc fusion protein, the onset of arthritic symptoms were delayed by approximately 10 days as compared to the PBS treated mice. This demonstrates that the inhibition of the CRP-1/B7RP-1 pathway can alleviate disease symptoms in this mouse model of rheumatoid arthritis.

15 Mice treated with B7RP-1-Fc or B7.2-Fc showed an earlier onset of disease (Table 1 and Figure 16a) with an increase in arthritic severity (Figure 16b) as compared to the PBS-treated controls. This indicates that the B7RP-1-Fc fusion protein enhances the T-cell
20 immune response. Such activity may be useful in generating anti-tumor immunity *in vivo*.

The opposite effects by CRP-1-Fc and B7RP-1-Fc in this mouse model of rheumatoid arthritis indicate that the pathway can be manipulated to either enhance
25 or inhibit the disease progression. Targeting the CRP-1 protein with soluble B7RP-1-Fc enhances the disease, while the interaction of soluble CRP-1-Fc with B7RP-1 inhibits the disease symptoms.

30

Example 19

B7RP-1-Fc induces an inflammatory bowel disease phenotype in transgenic mice.

35

Persistent overexpression of the B7-related protein (B7RP1-Fc) in 22-to-25-week-old transgenic mice(Example 12) induces a striking phenotype of

- 99 -

inflammatory bowel disease (IBD) with marked thickening and chronic inflammation of the small and large bowels (enterocolitis) and weight loss in some animals. Histologically, the most severe inflammatory changes were found in the proximal and distal colon, with milder changes in the small intestine. The proximal colon was markedly thickened with fissuring ulceration, transmural inflammation, and hypertrophy of the colonic mucosa, while the distal colon had diffuse mucosal hypertrophy (or focal erosion and glandular atrophy) without ulceration. The proximal small intestine had mild to marked mucosal hypertrophy with milder inflammatory changes, while the distal small intestine (ileum) had mild mucosal hypertrophy in some animals and atrophy in other mice. The intestinal changes were most severe and consistently found in the female B7RP-1 transgenic mice, but were also observed in several of the male transgenic mice in this study.

It is interesting to note that the histologic features found in the proximal colon, including the fissuring ulceration and the transmural chronic granulomatous inflammation with multinucleated giant cells, more closely resemble those seen in Crohn's disease than ulcerative colitis in humans. Morphologically, this colitis also mimics the IBD described in mice deficient in interleukin-10, which develop wasting, anemia, and enterocolitis affecting their entire intestinal tract (Kuhn et al. 1993; Sartor 1995; Leach et al. 1999). As in the IL-10 knockout mice, the initial changes in the B7RP-1-Fc transgenic mice consist of mild, focal infiltrates of inflammatory cells in the lamina propria without colonic epithelial hyperplasia (Example 13). In older mice, the affected colonic segments become thickened due to glandular hypertrophy/hyperplasia and chronic inflammation.

- 100 -

The proximal and distal colons of the B7RP-1-Fc mice had moderate to severe colitis with histologic features of inflammatory bowel disease (IBD). The affected segments of the proximal colon (Figure 17B-17D) were
5 diffusely thickened, due to prominent glandular epithelial hypertrophy and hyperplasia with elongation and dilatation of the mucosal glands (Fig. 17B), which had increased numbers of mitotic figures and rare crypt abscesses, but retained goblet cells with mucin (Fig.
10 17D). The mucosa had diffuse chronic inflammation in the lamina propria, which in some animals extended transmurally to involve the underlying layers of the gut wall, including the submucosa, muscularis, serosa, and the adjacent mesenteric fat tissue (Fig. 17B-17C).
15 The inflammatory infiltrates consisted of lymphocytes (predominantly CD3+, CD44+ T cells), plasma cells, and epithelioid macrophages (Fig. 17F) mixed with some neutrophils and occasional multinucleated giant cells (Fig. 17E), characteristic of chronic granulomatous
20 inflammation. Lymphoid aggregates (mostly B220+ cells mixed with small numbers of CD3+ cells) were also present in the mucosa and around smaller blood vessels in the submucosa and deeper layers, including mesenteric fat (Fig. 17C). The lumen contained
25 mucopurulent or mucous exudate (Fig. 17D). Severe evidence of colitis, with multifocal fissuring ulceration of the mucosa and transmural inflammation (Fig. 17B-17C), was found in these B7RP-1-Fc transgenic mice.

30 The distal colon of the B7RP-1-Fc transgenic mice was also diffusely thickened and hyperplastic with elongation, basophilia, and dilatation of the colonic glands (Fig. 18B-18G), some of which contained crypt abscesses (Fig. 18D and 18F) and mucus. The lamina
35 propria had a mild diffuse inflammatory infiltrate of lymphocytes (predominantly CD3+, CD44+ cells,

- 101 -

particularly in the superficial mucosa; Fig. 18E), as well as plasma cells and focal aggregates of epithelioid macrophages mixed with some neutrophils. Lymphoid aggregates (of predominantly B220+ cells; Fig. 18D and 18F) were also scattered throughout the mucosa. The small intestine of B7RP-1-Fc transgenic mice had more variable changes, including mild to focally marked mucosal and crypt hypertrophy and hyperplasia (Fig. 19B and 19D with crypt/villus ratios ranging from 1:4 to 1.5:1, as compared to 1:10 in the control mice) accompanied by a predominantly lymphoplasmacytic infiltrate in the lamina propria. The mucosal hyperplasia was most prominent in the proximal small intestine, including the duodenum (Fig. 19B) and particularly the jejunum (Fig. 19D). The crypt architecture was focally deranged and dysplastic in the most severely affected mice (Fig. 19D). In contrast, the distal small intestines (ileum) of some mice, had mild, patchy villous atrophy of the ileal mucosa (Fig. 19F) with blunting, thickening or focal loss of villi (with a crypt:villus ratio of 1:1 or less, instead of the normal ratio of 1:2), while other mice had mild ileal mucosal hypertrophy.

The B7RP-1-Fc fusion protein acts to activate cells that are responsible for eliciting a phenotype very similar to that of human Crohn's disease. This indicates that the cells that may be responsible for the inflammation in Crohn's disease are activated by the B7RP-1-Fc fusion protein. Soluble protein, antibody or small molecule inhibitors of B7RP-1 may therefore be useful in inhibiting IBD.

Example 20

The B7RP-1-Fc fusion protein inhibits tumor growth in mice

- 102 -

To examine the effect of B7RP-1 and CRP-1 on the growth of the immunogenic murine Meth A sarcoma, we investigated whether the soluble B7RP-1-Fc affects the growth of an established Meth A sarcoma in Balb/c mice.

Exponentially growing Meth A sarcoma cells were implanted by intradermal injection of 0.5 million cells in the abdomen of Balb/c mice on day 0. On day 7, when the tumors reached ~ 100 mm³, the mice were treated with either vehicle (PBS) or B7RP-1-Fc (8mg/kg), subcutaneously in the neck on days 7, 10, 14, and 17. The bidimensional diameters of the tumors were measured by calipers and the tumor volume (in mm³) was estimated using the formula: Tumor volume = [(width)² × length] / 2. The tumor growth was monitored up to day 28. Each group had eight mice.

The Meth A sarcoma growth pattern of the control tumor was bi-phasic: a slow initial phase was followed by a relatively rapid exponential phase. In B7RP-1-Fc treated mice, the growth of the tumor was significantly slower in the rapid exponential phase. On day 28, the average volumes of the control and B7RP1-Fc treated mice were 1410 mm³ and 580 mm³, respectively (Figure 20). Therefore, B7RP-1-Fc treatment inhibited tumor growth significantly in this model. The data strongly suggest the beneficial therapeutic utility of the soluble B7RP-1-Fc protein, and other activators of the B7RP-1/CRP-1 pathway, in the treatment of immunogenic tumors.

Immunologic anti-tumor activity is closely associated with cytolytic T-lymphocyte (CTL) function. Consistently, the B7RP-1-Fc protein is expressed on cytolytic CD8+ T-cells

- 103 -

(Example 9, Figure 6). These data strongly support B7RP-1 functions on cytolytic CD8+ T-cells. B7RP-1-Fc, or other stimulators of the B7RP-1/CRP-1 pathway, may therefore be used to enhance cytolytic T-cell and cellular immune functions for a number of non-cancer-related indications.

Example 21

10 Inhibition of human B7RP-1 activity in vitro

To determine if human B7RP-1 has positive co-stimulatory properties, we tested cells expressing human B7RP-1 and human B7RP-1-Fc fusion protein in T-cell proliferation assays. The human B7RP-1-Fc fusion protein was constructed by fusing gene sequences corresponding to amino acids 1 to 247 to a partial human IgG1 gene sequence (Example 14). The human CRP-1-Fc fusion protein was constructed by fusing gene sequences corresponding to amino acids 1 to 146 to a partial human IgG1 gene sequence (Example 2). The methods of construction, expression and purification of both fusion proteins were conducted as described in Example 7. B7RP-1-Fc demonstrated co-stimulatory activities that are dependent on anti-CD3 stimulation (Figure 21a). In addition, this activity can be specifically inhibited with soluble CRP-1-Fc protein (Figure 21b). Similar co-stimulatory effects were obtained using CHO cells that express membrane-bound, human B7RP-1, containing the entire coding sequence (Figure 21c).

The production of cytokines by human T-cells under the above *in vitro* proliferation conditions was determined. Supernatants from T-cell cultures stimulated for 48 and 72 hours were

- 104 -

analyzed for IL-2, IL-10, and IFN-gamma by ELISA according to the manufacturer's specifications (BioSource International). The IFN-gamma and IL-10 levels were significantly increased; however, unlike the case with CD28 co-stimulation, IL-2 was not notably induced (Figure 21d). The increased levels of IFN-gamma, a Th1 cytokine, correlate with the B7RP-1 functions to increase IgG2a, as described in Example 13.

10 *In vitro* T-cell co-stimulation assays were conducted as follows. Highly purified human T-cells (>98% CD3+) were isolated by negative selection of fresh or thawed, adherence depleted PBMC using mAb labeled magnetic beads (Miltenyi Biotec). T-cells (1×10^5 cells/well) were cultured in triplicate wells in 96 well plates in 200 μ l/well RPMI + 10% FCS. To evaluate B7RP-1-Fc co-stimulation, various concentrations of anti-CD3 (Pharmingen) and 10 μ g/ml anti-human IgG Fc (Sigma) in 100 μ l 1X PBS were pre-coated onto U bottom plates by an incubation at 4°C overnight. The unbound anti-CD3 and anti-human IgG Fc were removed, and the cells were cultured in the presence or absence of various concentrations of B7RP-1-Fc, OPG-Fc control or anti-CD28 (Pharmingen). For CRP-1-Fc inhibition of B7RP-1-Fc co-stimulation, T-cells were cultured in 0.33 μ g/ml anti-CD3 and 10 μ g/ml anti-human IgG Fc pre-coated wells with 0.5 μ g/ml B7RP-1-Fc in the presence of serially diluted CRP-1-Fc or OPG-Fc, starting at 10 μ g/ml. To evaluate co-stimulation by CHO cells expressing B7RP-1, T-cells were cultured in flat bottom plates with various concentrations of soluble anti-CD3 in the presence or absence of

- 105 -

various amounts of mitomycin-C treated CHO B7RP-1 cells or CHO vector cells. To test for T-cell proliferation, cultures were pulsed with 1 uCi/well [³H]TdR during the last 18 hrs of a 72 hr culture.

5 T-cell proliferation was determined by [³H]TdR incorporation. The results of one representative experiment from three random donors are expressed as mean CPM incorporated +/- SD. For analyses of cytokine production, cells were cultured for 48 and
10 72 hours and supernatants were collected for ELISA.

These experiments show that the extracellular portion of human B7RP-1, as described in Example 14, when fused to a human Fc fragment, can co-stimulate T-cells in vitro. This co-
15 stimulation is inhibited by CRP-1-Fc and thus demonstrates how a soluble inhibitor of human B7RP-1 may function. In vitro assays, such as that described here using human B7RP-1 and CRP-1, could be used to screen for antibody, soluble protein,
20 peptibody, or small molecule inhibitors of B7RP-1/CRP-1 activity.

* * *

25 While the present invention has been described in terms of the preferred embodiments, it is understood that variations and modifications will occur to those skilled in the art. Therefore, it is intended that the appended claims cover all such equivalent
30 variations that come within the scope of the invention as claimed.

- 106 -

WHAT IS CLAIMED IS:

1. An isolated nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:
- a) the nucleotide sequence as set forth in Figure 1A (SEQ ID NO: 1);
 - b) the nucleotide sequence encoding the polypeptide from residues 1-200 or from residues 21-200 as set forth in Figure 1A (SEQ ID NO: 1);
 - c) a nucleotide sequence encoding a polypeptide that is at least about 70 percent identical to the polypeptide as set forth in Figure 1A (SEQ ID NO: 1);
 - d) a naturally occurring allelic variant or alternate splice variant of any of (a), (b) or (c);
 - e) a nucleotide sequence complementary to any of (a), (b) or (c);
 - f) a nucleotide sequence of (b), (c) or (d) encoding a polypeptide fragment of at least about 25, 50, 75, 100, or greater than 100 amino acid residues;
 - g) a nucleotide sequence of (a), (b) or (c) comprising a fragment of at least about 10, 15, 20, 25, 50, 75, 100, or greater than 100 nucleotides; and
 - h) a nucleotide sequence which hybridizes under stringent conditions to any of (a)-(g).

2. An isolated nucleic acid molecule comprising a nucleotide sequence selected from the group consisting of:
- a) the nucleotide sequence as set forth in Figure 2A (SEQ ID NO: 11) or Figure 3A (SEQ ID NO: 6) or Figure 12A (SEQ ID NO: 16);
 - b) the nucleotide sequence encoding the polypeptide as set forth in Figure 2A (SEQ ID NO: 6) from residues 1-322 or from residues 47-322, or as set

- 107 -

forth in Figure 3A (SEQ ID NO: 11) from residues 1-288
or from residues 19-288, 20-288, 21-288, 22-288, 24-
288, or 28-288 or as set forth in Figure 12A from
residues 1-302, or from residues 19-302, 20-302, 21-
5 302, 22-302, 24-302 or 28-302;

c) a nucleotide sequence encoding a
polypeptide that is at least about 70 percent identical
to the polypeptide as set forth in Figure 2A (SEQ ID
NO: 6) or Figure 3A (SEQ ID NO: 11) or Figure 12A (SEQ
10 ID NO: 6);

d) a naturally occurring allelic variant or
alternate splice variant of any of (a), (b) or (c);

e) a nucleotide sequence complementary to any
of (a), (b) or (c);

15 f) a nucleotide sequence of (b), (c) or (d)
encoding a polypeptide fragment of at least about 25,
50, 75, 100, or greater than 100 amino acid residues;

g) a nucleotide sequence of (a), (b) or (c)
comprising a fragment of at least about 10, 15, 20, 25,
20 50, 75, 100, or greater than 100 nucleotides; and

h) a nucleotide sequence which hybridizes
under stringent conditions to any of (a)-(g).

3. The nucleic acid molecule of Claims 1 or 2
25 wherein the nucleotide sequence is operably linked to
an expression control sequence.

4. A host cell comprising the nucleic acid
molecule of Claim 2.

30

5. The host cell of Claim 3 which is a eucaryotic
cell.

6. The host cell of Claim 3 which is a procaryotic
35 cell.

- 108 -

7. A process for producing a polypeptide comprising growing a culture of the host cell of Claim 3 in suitable culture medium and isolating the polypeptide from the culture.

5

8. A polypeptide produced by the process of Claim 7.

9. A polypeptide encoded by the nucleic acid molecule of Claim 1.

10. A polypeptide encoded by the nucleic acid molecule of Claim 2.

11. An isolated polypeptide comprising the amino acid sequence selected from the group consisting of:

a) the amino acid sequence as set forth in Figure 1A (SEQ ID NO: 2);

b) the mature amino acid sequence as set forth in Figure 1A (SEQ ID NO: 2) comprising a mature amino terminus at residue 21;

c) a fragment of the amino acid sequence set forth in Figure 1A (SEQ ID NO: 2) comprising at least about 25, 50, 75, 100, or greater than 100 amino acid residues;

d) an ortholog of (a), (b) or (c); and

e) an allelic variant or alternative splice variant of (a), (b), (c) or (d).

12. An isolated polypeptide comprising the amino acid sequence selected from the group consisting of:

a) the amino acid sequence as set forth in Figure 2A (SEQ ID NO: 7) or Figure 3A (SEQ ID NO: 12) or Figure 12A (SEQ ID NO: 17);

b) the mature amino acid sequence as set forth in Figure 2A (SEQ ID NO: 7) comprising a mature

- 109 -

amino terminus at residues 47, or Figure 3A (SEQ ID NO: 12) comprising a mature amino terminus at any of residues 19, 20, 21, 22, 24 or 28, or Figure 12A (SEQ ID NO: 17) comprising a mature amino terminus at any of
5 residues 19,20,21,22,24,or 28;

c) a fragment of the amino acid sequence set forth in Figure 2A (SEQ ID NO: 7) or Figure 3A (SEQ ID NO: 12) or Figure 12A (SEQ ID NO: 17) comprising at least about 25, 50, 75, 100, or greater than 100 amino
10 acid residues;

d) an ortholog of (a), (b) or (c); and

e) an allelic variant or alternative splice variant of (a), (b), (c) or (d).

15 13. An antibody or fragment thereof specifically binding the polypeptide of Claims 9, 10, 11 or 12.

14. The antibody of Claim 11 which is a monoclonal antibody.

20

15. The antibody of Claim 13 which is a human antibody.

16. The antibody of Claim 13 which is a humanized
25 or CDR-grafted antibody.

17. The antibody or fragment of Claim 13 which binds B7RP1 or to a B7RP1 extracellular domain.

30 18. The antibody or fragment of Claim 13 which inhibits the binding of B7RP1 to CRP1.

19. A composition comprising the polypeptide of Claims 9, 10, 11 or 12 and a pharmaceutically
35 acceptable carrier, adjuvant, solubilizer, stabilizer or anti-oxidant.

- 110 -

20. A polypeptide comprising a derivative of the polypeptide of Claims 9, 10, 11 or 12.

5 21. The polypeptide of Claim 20 which is covalently modified with a water-soluble polymer.

22. A fusion polypeptide comprising the polypeptide of Claims 9, 10, 11 or 12 fused to a
10 heterologous amino acid sequence.

23. The fusion polypeptide of Claim 22 wherein the heterologous amino acid sequence is an IgG constant domain or fragment thereof.

15 24. A method for treating, preventing or ameliorating a T cell mediated disorder comprising administering to an animal the polypeptide of Claims 9, 10, 11 or 12.

20 25. A method of diagnosing a T cell mediated disorder or a susceptibility to a T cell mediated disorder in an animal comprising:

25 a) determining the presence or amount of expression of the polypeptide of Claims 9, 10, 11 or 12; and

30 b) diagnosing a T-cell mediated disorder or a susceptibility to a T-cell mediated disorder based on the presence or amount of expression of the polypeptide.

26. A method of identifying a test molecule which binds to a polypeptide comprising:

35 a) contacting the polypeptide of Claims 9, 10, 11 or 12 with a test molecule; and

- 111 -

b) determining the extent of binding of the polypeptide to the test molecule.

27. The method of Claim 26 further comprising
5 determining the activity of the polypeptide when bound to the compound.

28. A method of regulating T cell activation or proliferation in an animal comprising administering to
10 the animal the nucleic acid molecule of Claims 1, 2 or 3.

29. A transgenic non-human mammal comprising the nucleic acid molecule of Claim 3.
15

30. A method of suppressing an immune response in an animal comprising administering to the animal an antagonist of CRP-1 or B7RP-1.

20 31. The method of Claim 30 wherein the antagonist is an antibody which binds B7RP-1.

FIGURE 1A

ATG AAG CCG TAC TTC TGC CGT GTC TTT GTC TTC TGC TTC CTA ATC	45
M K P Y F C R V F V F C F L I	
5 10 15	
AGA CTT TTA ACA GGA GAA ATC AAT GGC TCG GCC GAT CAT AGG ATG	90
R L L T G E I N G S A D H R M	
20 25 30	
TTT TCA TTT CAC AAT GGA GGT GTA CAG ATT TCT TGT AAA TAC CCT	135
F S F H N G G V Q I S C K Y P	
35 40 45	
GAG ACT GTC CAG CAG TTA AAA ATG CGA TTG TTC AGA GAG AGA GAA	180
E T V Q Q L K M R L F R E R E	
50 55 60	
GTC CTC TGC GAA CTC ACC AAG ACC AAG GGA AGC GGA AAT GCG GTG	225
V L C E L T K T K G S G N A V	
65 70 75	
TCC ATC AAG AAT CCA ATG CTC TGT CTA TAT CAT CTG TCA AAC AAC	270
S I K N P M L C L Y H L S N N	
80 85 90	
AGC GTC TCT TTT TTC CTA AAC AAC CCA GAC AGC TCC CAG GGA AGC	315
S V S F F L N N P D S S Q G S	
95 100 105	
TAT TAC TTC TGC AGC CTG TCC ATT TTT GAC CCA CCT CCT TTT CAA	360
Y Y F C S L S I F D P P P F Q	
110 115 120	
GAA AGG AAC CTT AGT GGA GGA TAT TTG CAT ATT TAT GAA TCC CAG	405
E R N L S G G Y L H I Y E S Q	
125 130 135	
CTC TGC TGC CAG CTG AAG CTC TGG CTA CCC GTA GGG TGT GCA GCT	450
L C C Q L K L W L P V G C A A	
140 145 150	
TTC GTT GTG GTA CTC CTT TTT GGA TGC ATA CTT ATC ATC TGG TTT	495
F V V V L L F G C I L I I W F	
155 160 165	
TCA AAA AAG AAA TAC GGA TCC AGT GTG CAT GAC CCT AAT AGT GAA	540
S K K K Y G S S V H D P N S E	
170 175 180	
TAC ATG TTC ATG GCG GCA GTC AAC ACA AAC AAA AAG TCT AGA CTT	585
Y M F M A A V N T N K K S R L	
185 190 195	
GCA GGT GTG ACC TCA	600
A G V T S	
200	

FIGURE 1B

mCRP1	MKPYFCRVFV	FCFLIRLL--	-----TGEIN	GS----	ADHR	MFSFHNGGVQ	39
mCD28	MT-----	----LRLFL	ALNFFSVQVT	ENKILVKQSP	LLVVDSEVS		38
Consensus	M.....RL..V.		
mCRP1	ISCKYPETV-	-QQLKMRLFR	--EREV-LCE	LTKTKGSGNA	VSIKNPMLCL		34
mCD28	LSCRYSYNLL	AKEFRASLYK	GVNSDVEVCV	GNGNFTYQPQ	FRSNAEFNCD		38
Consensus	.SC.Y.....L..V..C.C.		
mCRP1	YHLSNNSVSF	FLNNPDSSQG	SYFCSLSIF	DPPPFQERNL	SGGYL-HIYE		133
mCD28	GDFDNETVTF	RLWNLHVNHT	DIYFCKIEFM	YPPPYLDNER	SNGTIIHIKE		138
Consensus	...N..V.F	.L.N.....	..YFC.....	.PPP.....	S.G...HI.E		
mCRP1	SQLC---CQL	KL-W-LPVGC	AA-FVVLLF	GCIL-IIWFS	KKKY----GS		172
mCD28	KHLCHTQSSP	KLFWALVVA	GVLFCYGLLV	TVALCVIWTN	SRRNRLQVT		188
Consensus	..LC.....	KL.W.L.V..	...F...LL.	...L..IW..		
mCRP1	SVH-DPNSEY	MFMAAVNTNK	KSR-LAGVTS				200
mCD28	TMMTERRPG	LTRKPYQPYA	PARDFAAYRP				218
ConsensusP....R..A....	...			

FIGURE 2A

ATG CAG CTA AAG TGT CCC TGT TTT GTG TCC TTG GGA ACC AGG CAG	45
M Q L K C P C F V S L G T R Q	5 10 15
CCT GTT TGG AAG AAG CTC CAT GTT TCT AGC GGG TTC TTT TCT GGT	90
P V W K K L H V S S G F F S G	20 25 30
CTT GGT CTG TTC TTG CTG CTG TTG AGC AGC CTC TGT GCT GCC TCT	135
L G L F L L L L S S L C A A S	35 40 45
GCA GAG ACT GAA GTC GGT GCA ATG GTG GGC AGC AAT GTG GTG CTC	180
A *E T E V G A M V G S N V V L	50 55 60
AGC TGC ATT GAC CCC CAC AGA CGC CAT TTC AAC TTG AGT GGT CTG	225
S C I D P H R R H F N L S G L	65 70 75
TAT GTC TAT TGG CAA ATC GAA AAC CCA GAA GTT TCG GTG ACT TAC	270
Y V Y W Q I E N P E V S V T Y	80 85 90
TAC CTG CCT TAC AAG TCT CCA GGG ATC AAT GTG GAC AGT TCC TAC	315
Y L P Y K S P G I N V D S S Y	95 100 105
AAG AAC AGG GGC CAT CTG TCC CTG GAC TCC ATG AAG CAG GGT AAC	360
K N R G H L S L D S M K Q G N	110 115 120
TTC TCT CTG TAC CTG AAG AAT GTC ACC CCT CAG GAT ACC CAG GAG	405
F S L Y L K N V T P Q D T Q E	125 130 135
TTC ACA TGC CGG GTA TTT ATG AAT ACA GCC ACA GAG TTA GTC AAG	450
F T C R V F M N T A T E L V K	140 145 150
ATC TTG GAA GAG GTG GTC AGG CTG CGT GTG GCA GCA AAC TTC AGT	495
I L E E V V R L R V A A N F S	155 160 165
ACA CCT GTC ATC AGC ACC TCT GAT AGC TCC AAC CCG GGC CAG GAA	540
T P V I S T S D S S N P G Q E	170 175 180
CGT ACC TAC ACC TGC ATG TCC AAG AAT GGC TAC CCA GAG CCC AAC	585
R T Y T C M S K N G Y P E P N	185 190 195
CTG TAT TGG ATC AAC ACA ACG GAC AAT AGC CTA ATA GAC ACG GCT	630
L Y W I N T T D N S L I D T A	200 205 210
CTG CAG AAT AAC ACT GTC TAC TTG AAC AAG TTG GGC CTG TAT GAT	675
L Q N N T V Y L N K L G L Y D	215 220 225
GTA ATC AGC ACA TTA AGG CTC CCT TGG ACA TCT CGT GGG GAT GTT	720
V I S T L R L P W T S R G D V	230 235 240

FIGURE 2A (Con't)

CTG TGC TGC GTA GAG AAT GTG GCT CTC CAC CAG AAC ATC ACT AGC	755
L C C V E N V A L H Q N I T S	
245 250 255	
ATT AGC CAG GCA GAA AGT TTC ACT GGA AAT AAC ACA AAG AAC CCA	810
I S Q A E S F T G N N T K N P	
260 265 270	
CAG GAA ACC CAC AAT AAT GAG TTA AAA GTC CTT GTC CCC GTC CTT	855
Q E T H N N E L K V L V P V L	
275 280 285	
GCT GTA CTG GCG GCA GCG GCA TTC GTT TCC TTC ATC ATA TAC AGA	900
A V L A A A A F V S F I I Y R	
290 295 300	
CGC ACG CGT CCC CAC CGA AGC TAT ACA GGA CCC AAG ACT GTA CAG	945
R T R P H R S Y T G P K T V Q	
305 310 315	
CTT GAA CTT ACA GAC CAC GCC	966
L E L T D H A	
320 322	

FIGURE 2B

mB7RP1	MQLKPCFVS	LGTRQPVWKK	LHVSSGFFSG	LGLFLLLLS-	SLCAASAETE	43
mCD80	MA--CNC--Q	LMQDTPL---	LKFPCPRI-	L-LFVLLIRL	SQVSSDVDEQ	41
Consensus	M...C.C...	L....P....	L.....	L.LF.LL...	S.....	
mB7RP1	VGAMVGSNVV	LSCIDPHRRH	FNLGLYVYW	QIENPEVSVT	YYLPYKSPGI	99
mCD80	LSKSVKDKVL	LPC-RYNPH	EDESEDRIYW	QKHDKV---	--LSVIAGKL	35
ConsensusV...V.	L.C....H	...S....YW	Q...V...	..L.....	
mB7RP1	NVDSSYKNRG	HLSLDSMKQG	NFSLYLKNVT	PQDTQEFTCR	VFMNTATELV	149
mCD80	KVWPEYKNR-	--TL--YDNT	TYSLIILGLV	LSDRGTYSKV	VQKKERGTYE	130
Consensus	.V...YKNR.	...L.....	..SL.....	..D.....C.	V.....	
mB7RP1	KILEEVRLR	VAANFSTPVI	STSDSSNPGQ	ERTYTCMSKN	GYPEPNLYWI	199
mCD80	VKHLALVKLS	IKADFSTPNI	TESGNPSADT	KRI-TCFASG	GFPKPRFSWL	179
ConsensusV.L.	...A.FSTP.I	..S.....	.R..TC...	G.P.P...W.	
mB7RP1	-NTTDSLID	TALQNTTVYL	NKLGLYDVIS	TLRLPWTSRG	DVLCCVENVA	248
mCD80	ENGRELPGIN	TTISQDPESE	LYTISSQLDF	NTTRNHTIKC	LIKYGDAHVS	229
Consensus	.N.....I.	T.....T...V.	
mB7RP1	LHQNITSISQ	AESFTGNNTK	NPQETHNNEL	KVLVPVLAVL	A-AAAFVSFI	237
mCD80	EDFTWEKPPE	DPPDSKNTLV	LFGAGFGAVI	TVVIVVVIK	CFCKHRSCFR	279
ConsensusN...V.V.V.....F.	
mB7RP1	IYRRTR-PHR	SYT-GPKTVQ	LELTDHA			322
mCD80	RNEASRETNN	SLTFGPPEAL	AEQTVFL			306
ConsensusR....	S.T.GP...	.E.T...			

FIGURE 3A

ATG CGG CTG GGC AGT CCT GGA CTG CTC TTC CTG CTC TTC AGC AGC	45
M R L G S P G L L F L L F S S	
5 10 15	
CTT CGA GCT GAT ACT CAG GAG AAG GAA GTC AGA GCG ATG GTA GGC	90
L R A *D *T *Q *E K *E V R A *M V G	
20 25 30	
AGC GAC GTG GAG CTC AGC TGC GCT TGC CCT GAA GGA AGC CGT TTT	135
S D V E L S C A C P E G S R F	
35 40 45	
GAT TTA AAT GAT GTT TAC GTA TAT TGG CAA ACC AGT GAG TCG AAA	180
D L N D V Y V Y W Q T S E S K	
50 55 60	
ACC GTG GTG ACC TAC CAC ATC CCA CAG AAC AGC TCC TTG GAA AAC	225
T V V T Y H I P Q N S S L E N	
65 70 75	
GTG GAC AGC CGC TAC CGG AAC CGA GCC CTG ATG TCA CCG GCC GGC	270
V D S R Y R N R A L M S P A G	
80 85 90	
ATG CTG CGG GGC GAC TTC TCC CTG CGC TTG TTC AAC GTC ACC CCC	315
M L R G D F S L R L F N V T P	
95 100 105	
CAG GAC GAG CAG AAG TTT CAC TGC CTG GTG TTG AGC CAA TCC CTG	360
Q D E Q K F H C L V L S Q S L	
110 115 120	
GGA TTC CAG GAG GTT TTG AGC GTT GAG GTT ACA CTG CAT GTG GCA	405
G F Q E V L S V E V T L H V A	
125 130 135	
GCA AAC TTC AGC GTG CCC GTC GTC AGC GCC CCC CAC AGC CCC TCC	450
A N F S V P V V S A P H S P S	
140 145 150	
CAG GAT GAG CTC ACC TTC ACG TGT ACA TCC ATA AAC GGC TAC CCC	495
Q D E L T F T C T S I N G Y P	
155 160 165	
AGG CCC AAC GTG TAC TGG ATC AAT AAG ACG GAC AAC AGC CTG CTG	540
R P N V Y W I N K T D N S L L	
170 175 180	
GAC CAG GCT CTG CAG AAT GAC ACC GTC TTC TTG AAC ATG CGG GGC	585
D Q A L Q N D T V F L N M R G	
185 190 195	
TTG TAT GAC GTG GTC AGC GTG CTG AGG ATC GCA CGG ACC CCC AGC	630
L Y D V V S V L R I A R T P S	
200 205 210	
GTG AAC ATT GGC TGC TGC ATA GAG AAC GTG CTT CTG CAG CAG AAC	675
V N I G C C I E N V L L Q Q N	
215 220 225	
CTG ACT GTC GGC AGC CAG ACA GGA AAT GAC ATC GGA GAG AGA GAC	720
L T V G S Q T G N D I G E R D	
230 235 240	

7/33

FIGURE 3A (Con't)


[illegible]

FIGURE 3B

hB7RP1	EKEVRAMVGS	DVELSCACPE	GSRFDLNDVY	VYWQTSESKT	VVTYHIPQNS	50
mB7RP1	ETEVGAMVGS	NVVLSCIDPH	RRHFNLSGLY	VYWQIENPEV	SVTYYLBYKS	50
Consensus	E.EV.AMVGS	.V.LSC...P.	H...F.L...Y	VYWQ.....	.VTY...P..S	
hB7RP1	SLENVDSRYR	NRALMSPAGM	LRGDFSLRLF	NVTPQDEQKF	HCLVLSQ-SL	99
mB7RP1	PGINVDSSYK	NRGHLSLDSM	KQGNFSLYLK	NVTPQDTQEF	TCRVFMNTAT	100
Consensus	...NVDS.Y.	NR...S...M	..G.FSL.L.	NVTPQD.Q.F	.C.V.....	
hB7RP1	GFQEVLSVEV	TLHVAANFSV	PVVSAPHSPS	Q-DELTFTCT	SINGYPRPNV	143
mB7RP1	ELVKILEEVV	RLRVAANFST	PVISTSDSSN	PGQERTYTCM	SKNGYPEPNL	150
ConsensusL...V	.L.VAANFS.	PV.S...S..	...E.T.TC.	S.NGYP.PN.	
hB7RP1	YWINKTDNSL	LDQALQNDTV	FLNMRGLYDV	VSVLRIARTP	SVNIGCCCIEN	198
mB7RP1	YWINTTDNSL	IDTALQNNTV	YLNKLGLYDV	ISTLRLPWTS	RGDVLCCVEN	200
Consensus	YWIN.TDNSL	.D.ALQN.TV	.LN..GLYDV	.S.LR...T.CC.EN	
hB7RP1	VLLQQNLTVG	SQTGNDIGER	DKITENPVST	GEKNAATWSI	LAVLCLLVVV	248
mB7RP1	VALHQNITSI	SQAESFTGNN	TKNPQETHNN	ELKVLV--PV	LAVLAAAAFV	248
Consensus	V.L.QN.T..	SQ...F...G..	.K.....K...	..K...7...	LAVL.....V	
hB7RP1	AVAIGWVCRD	RCLQHSYAG				267
mB7RP1	SFIIYR--RT	R-PHRSYTGP	KTVQLELTDH	A		276
Consensus	...I...R.	R...SY.G.			

9/33

Figure 4A



10/33

Figure 4B





Figure 5



12/33

Figure 6






Figure 7



14/33

Figure 8



15/33

FIGURE 9




Figure 10




Figure 11




Figure 12A.

GCTGGTACGCCTGCAGGTACCGGTCCGGAATTC																CGGGTTCGACCCACGCGTCCGCCCACGCG																-138
TCCGCGGGAGCGCAGTTAGAGCCGATCTCCCGCGCCCCGAGGTTGCTCCTCTCCGAGGTCTC																																-76
CCGCGGCCCAAGTTCTCCGCGCCCCGAGGTCTCCGCGCCCCGAGGTCTCCGCGGCCCGAGGT																																-14
CTCCGCCCCGACC																																-1
ATG CGG CTG GGC AGT CCT GGA CTG CTC TTC CTG CTC TTC AGC AGC																45																
M R L G S P G L L F L L F S S																																
5 10 15																																
CTT CGA GCT GAT ACT CAG GAG AAG GAA GTC AGA GCG ATG GTA GGC																90																
L R A *D *T *Q *E K *E V R A *M V G																																
20 25 30																																
AGC GAC GTG GAG CTC AGC TGC GCT TGC CCT GAA GGA AGC CGT TTT																135																
S D V E L S C A C P E G S R F																																
35 40 45																																
GAT TTA AAT GAT GTT TAC GTA TAT TGG CAA ACC AGT GAG TCG AAA																180																
D L N D V Y V Y W Q T S E S K																																
50 55 60																																
ACC GTG GTG ACC TAC CAC ATC CCA CAG AAC AGC TCC TTG GAA AAC																225																
T V V T Y H I P Q N S S L E N																																
65 70 75																																
GTG GAC AGC CGC TAC CGG AAC CGA GCC CTG ATG TCA CCG GCC GGC																270																
V D S R Y R N R A L M S P A G																																
80 85 90																																
ATG CTG CGG GGC GAC TTC TCC CTG CGC TTG TTC AAC GTC ACC CCC																315																
M L R G D F S L R L F N V T P																																
95 100 105																																
CAG GAC GAG CAG AAG TTT CAC TGC CTG GTG TTG AGC CAA TCC CTG																360																
Q D E Q K F H C L V L S Q S L																																
110 115 120																																
GGA TTC CAG GAG GTT TTG AGC GTT GAG GTT ACA CTG CAT GTG GCA																405																
G F Q E V L S V E V T L H V A																																
125 130 135																																
GCA AAC TTC AGC GTG CCC GTC GTC AGC GCC CCC CAC AGC CCC TCC																450																
A N F S V P V V S A P H S P S																																
140 145 150																																
CAG GAT GAG CTC ACC TTC ACG TGT ACA TCC ATA AAC GGC TAC CCC																495																
Q D E L T F T C T S I N G Y P																																
155 160 165																																
AGG CCC AAC GTG TAC TGG ATC AAT AAG ACG GAC AAC AGC CTG CTG																540																
R P N V Y W I N K T D N S L L																																
170 175 180																																
GAC CAG GCT CTG CAG AAT GAC ACC GTC TTC TTG AAC ATG CGG GGC																585																
D Q A L Q N D T V F L N M R G																																
185 190 195																																
TTG TAT GAC GTG GTC AGC GTG CTG AGG ATC GCA CGG ACC CCC AGC																630																
L Y D V V S V L R I A R T P S																																
200 205 210																																
GTG AAC ATT GGC TGC TGC ATA GAG AAC GTG CTT CTG CAG CAG AAC																675																
V N I G C C I E N V L L Q Q N																																
215 220 225																																

19/33

CTG ACT GTC GGC AGC CAG ACA GGA AAT GAC ATC GGA GAG AGA GAC	720
L T V G S Q T G N D I G E R D	
230 235 240	
AAG ATC ACA GAG AAT CCA GTC AGT ACC GGC GAG AAA AAC GCG GCC	765
K I T E N P V S T G E K N A A	
245 250 255	
ACG TGG AGC ATC CTG GCT GTC CTG TGC CTG CTT GTG GTC GTG GCG	810
T W S I L A V L C L L V V V A	
260 265 270	
GTG GCC ATA GGC TGG GTG TGC AGG GAC CGA TGC CTC CAA CAC AGC	855
V A I G W V C R D R C L Q H S	
275 280 285	
TAT GCA GGT GCC TGG GCT GTG AGT CCG GAG ACA GAG CTC ACT GGC	900
Y A G A W A V S P E T E L T G	
300	
CAC GTT TGA	909
H V STOP	
302	
CCGGAGCTCACC GCCCAGAGCGTGGACAGGGCTTCCGTGAGACGCCACCGTGAGAGGCCAGG	971
TGGCAGCTTGAGCATGGACTCCCAGACTGCAGGGGAGCACTTGGGGCAGCCCCCAGAAGGAC	1033
CACTGCTGGATCCCAGGGAGAACCTGCTGGCGTTGGCTGTGATCCTGGAATGAGGCCCTTTC	1095

Figure 12B.

human	MRLGSP----	-----	-----G	L-LF-LLFSS	LRADTQEKEV	25
mouse	MQLKPCPCFVS	LGTRQPVWKK	LHVSSGFFSG	LGLFLLLLSS	LCAASAETEV	50
Consensus	M.L..P....G	L.LF.LL.SS	L.A...E.EV	50
human	RAMVGSDVEL	SCACPEGSRF	DLNDVYVYWQ	TSESKTVVVTY	HIPQNSSLN	75
mouse	GAMVGSNVVL	SCIDPHRRHF	NLSGLYVYWQ	IENPEVSVTY	YLPYKSPGIN	100
Consensus	.AMVGS.V.L	SC..P....F	.L...YVYWQVTY	..P..S...N	100
human	VDSRYRNRAL	MSPAGMLRGD	FSLRLFNVT	QDEQKFHCLV	LSQ-SLGFQE	124
mouse	VDSSYKNRGH	LSLDSMKQGN	FSLYLKNVT	QDTQEFTCRV	FMNTATELVK	150
Consensus	VDS.Y.NR..	.S...M..G.	FSL.L.NVT	QD.Q.F.C.V	150
human	VLSVEVTLHV	AANFSVPVVS	APHSPSQ-DE	LTFTCTSING	YPRPNVYWIN	173
mouse	ILEEVVRLRV	AANFSTPVIS	TSDSSNPGQE	RTYTCMSKNG	YPEPNLYWIN	200
Consensus	.L...V.L.V	AANFS.PV.S	...S.....E	.T.TC.S.NG	YP.PN.YWIN	200
human	KTDNSLLDQA	LQNDTVFLNM	RGLYDVVSVL	RIARTPSVNI	GCCIENVLLQ	223
mouse	TTDNSLIDTA	LQNNTVYLNK	LGLYDVISTL	RLPWTSRGDV	LCCVENVALH	250
Consensus	.TDNSL.D.A	LQN.TV.LN.	.GLYDV.S.L	R...T.....	.CC.ENV.L.	250
human	QNLTVGSQTG	NDIGERDKIT	ENPVSTGEKN	AATWSILAVL	CLLVVVAVAI	273
mouse	QNITSISQAE	SFTGNNTKNP	QETHNNELKV	LV--PVLAVL	AAAAFVSFII	298
Consensus	QN.T..SQ..	...G...K..K.LAVLV...I	300
human	GWVCRDRCLQ	HSYAGAWAVS	PETELTGHV			302
mouse	YR--RTR-PH	RSYTGPKTVQ	LE--LTDHA			322
ConsensusR.R...	.SY.G...V.	.E..LT.H.			329

Figure 13A

AACAATTTACACAGGAAACAGCTATGACCATGATTACGCCAAGCTCTAATACGA -111
 CTCACTATAGGGAAAGCTGGTACGCCTGCAGGTACCGGTCCGGAATTCCCGGGTC -56
 GACCCACGCGTCCGTGAACACTGAACGCGAGGACTGTTAAGTGTCTTGCAAAC -1

ATG AAG TCA GGC CTC TGG TAT TTC TTT CTC TTC TGC TTG CGC ATT 45
 M K S G L W Y F F L F C L R I
 5 10 15

AAA GTT TTA ACA GGA GAA ATC AAT GGT TCT GCC AAT TAT GAG ATG 90
 K V L T *G *E I N G S A N Y E M
 20 25 30

TTT ATA TTT CAC AAC GGA GGT GTA CAA ATT TTA TGC AAA TAT CCT 135
 F I F H N G G V Q I L C K Y P
 35 40 45

GAC ATT GTC CAG CAA TTT AAA ATG CAG TTG CTG AAA GGG GGG CAA 180
 D I V Q Q F K M Q L L K G G Q
 50 55 60

ATA CTC TGC GAT CTC ACT AAG ACA AAA GGA AGT GGA AAC ACA GTG 225
 I L C D L T K T K G S G N T V
 65 70 75

TCC ATT AAG AGT CTG AAA TTC TGC CAT TCT CAG TTA TCC AAC AAC 270
 S I K S L K F C H S Q L S N N
 80 85 90

AGT GTC TCT TTT TTT CTA TAC AAC TTG GAC CAT TCT CAT GCC AAC 315
 S V S F F L Y N L D H S H A N
 95 100 105

TAT TAC TTC TGC AAC CTA TCA ATT TTT GAT CCT CCT CCT TTT AAA 360
 Y Y F C N L S I F D P P P F K
 110 115 120

GTA ACT CTT ACA GGA GGA TAT TTG CAT ATT TAT GAA TCA CAA CTT 405
 V T L T G G Y L H I Y E S Q L
 125 130 135

TGT TGC CAG CTG AAG TTC TGG TTA CCC ATA GGA TGT GCA GCC TTT 450
 C C Q L K F W L P I G C A A F
 140 145 150

GTT GTA GTC TGC ATT TTG GGA TGC ATA CTT ATT TGT TGG CTT ACA 495
 V V V C I L G C I L I C W L T
 155 160 165

AAA AAG AAG TAT TCA TCC AGT GTG CAC GAC CCT AAC GGT GAA TAC 540
 K K K Y S S S V H D P N G E Y
 170 175 180

ATG TTC ATG AGA GCA GTG AAC ACA GCC AAA AAA TCT AGA CTC ACA 585
 M F M R A V N T A K K S R L T
 185 190 195

GAT GTG ACC CTA TAA 600
 D V T L STOP
 199

TATGGAAGTCTGGCAGCCAGGCATGAAGCACGTTGGCCAGTTTTCCTCAACTTGA 655
 AGTGCAAGATTCTCTTATTTCCGGGACCACGGAGAGTCTGACTTAACATACATA 710

TCTTCTGCTGGTGTGTTTTGTTCAATCTGGAAGAATGACTGTATCAGTCAATGGGGA	765
TTTTAACAGACTGCCTTGGTACTGCCGAGTCCTCTCAAAACAAACACCCCTCTTGC	820
AACCAGCTTTGGAGAAAGCCCAGCTCCTGTGTGCTCACTGGGAGTGGAATCCCTG	875
TCTCCACATCTGCTCCTAGCAGTGCATCAGCCAGTAAACAAACACATTTACAAG	930
AAAAATGTTTTAAAGATGCCAGGGGTACTGAATCTGCAAAGCAAATGAGCAGCCA	985
AGGACCAGCATCTGTCCGCATTTCACTATCATACTACCTCTTCTTTCTGTAGGGA	1040
TGAGAATTCCTCTTTTAATCAGTCAAGGGAGATGCTTCAAAGCTGGAGCTATTTT	1095
ATTTCTGAGATGTTGATGTGAAGTGTACATTAGTACATACTCAGTACTCTCCTTC	1150
AATTGCTGAACCCCAGTTGACCATTTTACCAAGACTTTAGATGCTTTCTTGTGCC	1205

Figure 13B

hCRP1	MKSGLWYFFLFCLRIKVLTEINGSANYEMFIFHNGGVQILCKYPDIVQQ	50
mCRP1	MKPYFCRVFVFCFLIRLLTGEINGSADHRMFSFHNGGVQISCKYPETVQQ	50
hCRP1	FKMQLLKGGQILCDLTKTKGSGNTVSIKSLKFCHSQLSNNSVSFFLYNLD	100
mCRP1	LKMRLFREREVLCELTKTKGSGNAVSIGNPMLCLYHLSNNSVSFFLNNPD	100
hCRP1	HSHANYYFCNLSIFDPPPFKV.TLTGGYLHIYESQLCCQLKFWLPIGCAA	149
mCRP1	SSQGSYYFCSLSIFDPPPFQERNLSGGYLHIYESQLCCQLKLWLPVGCAA	150
hCRP1	FVVVCILGCILICWLTKKKYSSSVHDPNGEYMFMRVNTAKKSRLTDVTL	199
mCRP1	FVVVLLFGCILIIFWSKKKYGSSVHDPNSEYMFMAAVNTNKSRLAGVTS	200




Figure 14





Figure 15

26/33

A.



B.




Figure 16

Figure 20




Figure 21A/B.





Figure 21C.






Figure 21D





A. Control mouse#53F: Prox. colon 40X



B. Mouse#111F: Prox. colon 40X




C. Mouse#111F: Prox. colon 20X



D. Mouse#111F: closeup of mucosa 100X




E. Mouse#112F: Giant cell, submucosa




F. Mouse#112F: epithelioid macrophages


Figure 17



A. Control mouse#53F:Distal colon, 40X



B. mouse#111F:Distal colitis, 40X



C. mouse#55M:Distal colitis, 40X




D. mouse#112F:Distal colon, 40X



E. mouse#112:CD3+ T-cells, 40X




F. mouse#112:closeup, 100X



A. Control mouse#53F:duodenum, 40X




B. Mouse#51F:duodenum, 40X




C. Control mouse#53F:jejunum, 40X



D. Mouse#51F:jejunal hyperplasia, 40X



E. Control mouse#53F:ileum, 40X



F. Mouse#231M:ileal atrophy, 40X

Figure 19

- 1 -

SEQUENCE LISTING

<110> AMGEN INC.

<120> NOVEL POLYPEPTIDES INVOLVED IN IMMUNE RESPONSE

<130> A-579A

<140> 09/264,527

<141> 1999-03-08

<160> 35

<170> PatentIn Ver. 2.1

<210> 1

<211> 600

<212> DNA

<213> mouse

<220>

<221> CDS

<222> Complement((1)..(600))

<400> 1

atg aag ccg tac ttc tgc cgt gtc ttt gtc ttc tgc ttc cta atc aga	48
Met Lys Pro Tyr Phe Cys Arg Val Phe Val Phe Cys Phe Leu Ile Arg	
1 5 10 15	
ctt tta aca gga gaa atc aat ggc tcg gcc gat cat agg atg ttt tca	96
Leu Leu Thr Gly Glu Ile Asn Gly Ser Ala Asp His Arg Met Phe Ser	
20 25 30	
ttt cac aat gga ggt gta cag att tct tgt aaa tac cct gag act gtc	144
Phe His Asn Gly Gly Val Gln Ile Ser Cys Lys Tyr Pro Glu Thr Val	
35 40 45	
cag cag tta aaa atg cga ttg ttc aga gag aga gaa gtc ctc tgc gaa	192
Gln Gln Leu Lys Met Arg Leu Phe Arg Glu Arg Glu Val Leu Cys Glu	
50 55 60	
ctc acc aag acc aag gga agc gga aat gcg gtg tcc atc aag aat cca	240
Leu Thr Lys Thr Lys Gly Ser Gly Asn Ala Val Ser Ile Lys Asn Pro	
65 70 75 80	
atg ctc tgt cta tat cat ctg tca aac aac agc gtc tct ttt ttc cta	288
Met Leu Cys Leu Tyr His Leu Ser Asn Asn Ser Val Ser Phe Phe Leu	
85 90 95	
aac aac cca gac agc tcc cag gga agc tat tac ttc tgc agc ctg tcc	336
Asn Asn Pro Asp Ser Ser Gln Gly Ser Tyr Tyr Phe Cys Ser Leu Ser	
100 105 110	
att ttt gac cca cct cct ttt caa gaa agg aac ctt agt gga gga tat	384
Ile Phe Asp Pro Pro Pro Phe Gln Glu Arg Asn Leu Ser Gly Gly Tyr	
115 120 125	
ttg cat att tat gaa tcc cag ctc tgc tgc cag ctg aag ctc tgg cta	432
Leu His Ile Tyr Glu Ser Gln Leu Cys Cys Gln Leu Lys Leu Trp Leu	
130 135 140	
ccc gta ggg tgt gca gct ttc gtt gtg gta ctc ctt ttt gga tgc ata	480
Pro Val Gly Cys Ala Ala Phe Val Val Val Leu Leu Phe Gly Cys Ile	
145 150 155 160	

ctt atc atc tgg ttt tca aaa aag aaa tac gga tcc agt gtg cat gac 528
 Leu Ile Ile Trp Phe Ser Lys Lys Lys Tyr Gly Ser Ser Val His Asp
 165 170 175
 cct aat agt gaa tac atg ttc atg gcg gca gtc aac aca aac aaa aag 576
 Pro Asn Ser Glu Tyr Met Phe Met Ala Ala Val Asn Thr Asn Lys Lys
 180 185 190
 tct aga ctt gca ggt gtg acc tca 600
 Ser Arg Leu Ala Gly Val Thr Ser
 195 200

<210> 2
 <211> 200
 <212> PRT
 <213> mouse

<400> 2
 Met Lys Pro Tyr Phe Cys Arg Val Phe Val Phe Cys Phe Leu Ile Arg
 1 5 10 15
 Leu Leu Thr Gly Glu Ile Asn Gly Ser Ala Asp His Arg Met Phe Ser
 20 25 30
 Phe His Asn Gly Gly Val Gln Ile Ser Cys Lys Tyr Pro Glu Thr Val
 35 40 45
 Gln Gln Leu Lys Met Arg Leu Phe Arg Glu Arg Glu Val Leu Cys Glu
 50 55 60
 Leu Thr Lys Thr Lys Gly Ser Gly Asn Ala Val Ser Ile Lys Asn Pro
 65 70 75 80
 Met Leu Cys Leu Tyr His Leu Ser Asn Asn Ser Val Ser Phe Phe Leu
 85 90 95
 Asn Asn Pro Asp Ser Ser Gln Gly Ser Tyr Tyr Phe Cys Ser Leu Ser
 100 105 110
 Ile Phe Asp Pro Pro Pro Phe Gln Glu Arg Asn Leu Ser Gly Gly Tyr
 115 120 125
 Leu His Ile Tyr Glu Ser Gln Leu Cys Cys Gln Leu Lys Leu Trp Leu
 130 135 140
 Pro Val Gly Cys Ala Ala Phe Val Val Val Leu Leu Phe Gly Cys Ile
 145 150 155 160
 Leu Ile Ile Trp Phe Ser Lys Lys Lys Tyr Gly Ser Ser Val His Asp
 165 170 175
 Pro Asn Ser Glu Tyr Met Phe Met Ala Ala Val Asn Thr Asn Lys Lys
 180 185 190
 Ser Arg Leu Ala Gly Val Thr Ser
 195 200

<210> 3
 <211> 200
 <212> PRT

<213> mouse

<400> 3

```

Met Lys Pro Tyr Phe Cys Arg Val Phe Val Phe Cys Phe Leu Ile Arg
 1          5          10          15

Leu Leu Thr Gly Glu Ile Asn Gly Ser Ala Asp His Arg Met Phe Ser
          20          25          30

Phe His Asn Gly Gly Val Gln Ile Ser Cys Lys Tyr Pro Glu Thr Val
          35          40          45

Gln Gln Leu Lys Met Arg Leu Phe Arg Glu Arg Glu Val Leu Cys Glu
          50          55          60

Leu Thr Lys Thr Lys Gly Ser Gly Asn Ala Val Ser Ile Lys Asn Pro
 65          70          75          80

Met Leu Cys Leu Tyr His Leu Ser Asn Asn Ser Val Ser Phe Phe Leu
          85          90          95

Asn Asn Pro Asp Ser Ser Gln Gly Ser Tyr Tyr Phe Cys Ser Leu Ser
          100          105          110

Ile Phe Asp Pro Pro Pro Phe Gln Glu Arg Asn Leu Ser Gly Gly Tyr
          115          120          125

Leu His Ile Tyr Glu Ser Gln Leu Cys Cys Gln Leu Lys Leu Trp Leu
          130          135          140

Pro Val Gly Cys Ala Ala Phe Val Val Val Leu Leu Phe Gly Cys Ile
          145          150          155          160

Leu Ile Ile Trp Phe Ser Lys Lys Lys Tyr Gly Ser Ser Val His Asp
          165          170          175

Pro Asn Ser Glu Tyr Met Phe Met Ala Ala Val Asn Thr Asn Lys Lys
          180          185          190

Ser Arg Leu Ala Gly Val Thr Ser
          195          200

```

<210> 4

<211> 218

<212> PRT

<213> mouse

<400> 4

```

Met Thr Leu Arg Leu Leu Phe Leu Ala Leu Asn Phe Phe Ser Val Gln
 1          5          10          15

Val Thr Glu Asn Lys Ile Leu Val Lys Gln Ser Pro Leu Leu Val Val
          20          25          30

Asp Ser Asn Glu Val Ser Leu Ser Cys Arg Tyr Ser Tyr Asn Leu Leu
          35          40          45

Ala Lys Glu Phe Arg Ala Ser Leu Tyr Lys Gly Val Asn Ser Asp Val
          50          55          60

Glu Val Cys Val Gly Asn Gly Asn Phe Thr Tyr Gln Pro Gln Phe Arg
          65          70          75          80

```

Ser Asn Ala Glu Phe Asn Cys Asp Gly Asp Phe Asp Asn Glu Thr Val
85 90 95

Thr Phe Arg Leu Trp Asn Leu His Val Asn His Thr Asp Ile Tyr Phe
100 105 110

Cys Lys Ile Glu Phe Met Tyr Pro Pro Pro Tyr Leu Asp Asn Glu Arg
115 120 125

Ser Asn Gly Thr Ile Ile His Ile Lys Glu Lys His Leu Cys His Thr
130 135 140

Gln Ser Ser Pro Lys Leu Phe Trp Ala Leu Val Val Val Ala Gly Val
145 150 155 160

Leu Phe Cys Tyr Gly Leu Leu Val Thr Val Ala Leu Cys Val Ile Trp
165 170 175

Thr Asn Ser Arg Arg Asn Arg Leu Leu Gln Val Thr Thr Met Asn Met
180 185 190

Thr Pro Arg Arg Pro Gly Leu Thr Arg Lys Pro Tyr Gln Pro Tyr Ala
195 200 205

Pro Ala Arg Asp Phe Ala Ala Tyr Arg Pro
210 215

<210> 5
<211> 234
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:Synthetic
Oglionucleotide

<400> 5
Met Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Arg
1 5 10 15

Leu Leu Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
20 25 30

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
35 40 45

Val Xaa Xaa Ser Cys Xaa Tyr Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
50 55 60

Xaa Xaa Xaa Leu Xaa Xaa Xaa Xaa Xaa Xaa Xaa Val Xaa Xaa Cys Xaa
65 70 75 80

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
85 90 95

Xaa Xaa Cys Xaa Xaa Xaa Xaa Asn Xaa Xaa Val Xaa Phe Xaa Leu
100 105 110

Xaa Asn Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Tyr Phe Cys Xaa Xaa Xaa
115 120 125

Xaa Xaa Xaa Pro Pro Pro Xaa Xaa Xaa Xaa Xaa Ser Xaa Gly Xaa
130 135 140

Xaa Xaa His Ile Xaa Glu Xaa Xaa Leu Cys Xaa Xaa Xaa Xaa Xaa Xaa
 145 150 155 160
 Lys Leu Xaa Trp Xaa Leu Xaa Val Xaa Xaa Xaa Xaa Phe Xaa Xaa
 165 170 175
 Xaa Xaa Leu Leu Xaa Xaa Xaa Xaa Leu Xaa Xaa Ile Trp Xaa Xaa Xaa
 180 185 190
 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Pro Xaa
 195 200 205
 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Arg
 210 215 220
 Xaa Xaa Ala Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
 225 230

<210> 6

<211> 966

<212> DNA

<213> mouse

<220>

<221> CDS

<222> Complement((1)..(966))

<400> 6

atg	cag	cta	aag	tgt	ccc	tgt	ttt	gtg	tcc	ttg	gga	acc	agg	cag	cct	48
Met	Gln	Leu	Lys	Cys	Pro	Cys	Phe	Val	Ser	Leu	Gly	Thr	Arg	Gln	Pro	
1				5					10					15		
ggt	tgg	aag	aag	ctc	cat	gtt	tct	agc	ggg	ttc	ttt	tct	ggt	ctt	ggt	96
Val	Trp	Lys	Lys	Leu	His	Val	Ser	Ser	Gly	Phe	Phe	Ser	Gly	Leu	Gly	
			20					25					30			
ctg	ttc	ttg	ctg	ctg	ttg	agc	agc	ctc	tgt	gct	gcc	tct	gca	gag	act	144
Leu	Phe	Leu	Leu	Leu	Leu	Ser	Ser	Leu	Cys	Ala	Ala	Ser	Ala	Glu	Thr	
			35				40					45				
gaa	gtc	ggt	gca	atg	gtg	ggc	agc	aat	gtg	gtg	ctc	agc	tgc	att	gac	192
Glu	Val	Gly	Ala	Met	Val	Gly	Ser	Asn	Val	Val	Leu	Ser	Cys	Ile	Asp	
	50					55					60					
ccc	cac	aga	cgc	cat	ttc	aac	ttg	agt	ggt	ctg	tat	gtc	tat	tgg	caa	240
Pro	His	Arg	Arg	His	Phe	Asn	Leu	Ser	Gly	Leu	Tyr	Val	Tyr	Trp	Gln	
					70					75					80	
atc	gaa	aac	cca	gaa	gtt	tcg	gtg	act	tac	tac	ctg	cct	tac	aag	tct	288
Ile	Glu	Asn	Pro	Glu	Val	Ser	Val	Thr	Tyr	Tyr	Leu	Pro	Tyr	Lys	Ser	
				85					90					95		
cca	ggg	atc	aat	gtg	gac	agt	tcc	tac	aag	aac	agg	ggc	cat	ctg	tcc	336
Pro	Gly	Ile	Asn	Val	Asp	Ser	Ser	Tyr	Lys	Asn	Arg	Gly	His	Leu	Ser	
			100					105					110			
ctg	gac	tcc	atg	aag	cag	ggt	aac	ttc	tct	ctg	tac	ctg	aag	aat	gtc	384
Leu	Asp	Ser	Met	Lys	Gln	Gly	Asn	Phe	Ser	Leu	Tyr	Leu	Lys	Asn	Val	
			115				120					125				
acc	cct	cag	gat	acc	cag	gag	ttc	aca	tgc	cgg	gta	ttt	atg	aat	aca	432
Thr	Pro	Gln	Asp	Thr	Gln	Glu	Phe	Thr	Cys	Arg	Val	Phe	Met	Asn	Thr	

130	135	140	
gcc aca gag tta gtc aag atc ttg gaa gag gtg gtc agg ctg cgt gtg Ala Thr Glu Leu Val Lys Ile Leu Glu Glu Val Val Arg Leu Arg Val 145 150 155 160			480
gca gca aac ttc agt aca cct gtc atc agc acc tct gat agc tcc aac Ala Ala Asn Phe Ser Thr Pro Val Ile Ser Thr Ser Asp Ser Ser Asn 165 170 175			528
ccg ggc cag gaa cgt acc tac acc tgc atg tcc aag aat ggc tac cca Pro Gly Gln Glu Arg Thr Tyr Thr Cys Met Ser Lys Asn Gly Tyr Pro 180 185 190			576
gag ccc aac ctg tat tgg atc aac aca acg gac aat agc cta ata gac Glu Pro Asn Leu Tyr Trp Ile Asn Thr Thr Asp Asn Ser Leu Ile Asp 195 200 205			624
acg gct ctg cag aat aac act gtc tac ttg aac aag ttg ggc ctg tat Thr Ala Leu Gln Asn Asn Thr Val Tyr Leu Asn Lys Leu Gly Leu Tyr 210 215 220			672
gat gta atc agc aca tta agg ctc cct tgg aca tct cgt ggg gat gtt Asp Val Ile Ser Thr Leu Arg Leu Pro Trp Thr Ser Arg Gly Asp Val 225 230 235 240			720
ctg tgc tgc gta gag aat gtg gct ctc cac cag aac atc act agc att Leu Cys Cys Val Glu Asn Val Ala Leu His Gln Asn Ile Thr Ser Ile 245 250 255			768
agc cag gca gaa agt ttc act gga aat aac aca aag aac cca cag gaa Ser Gln Ala Glu Ser Phe Thr Gly Asn Asn Thr Lys Asn Pro Gln Glu 260 265 270			816
acc cac aat aat gag tta aaa gtc ctt gtc ccc gtc ctt gct gta ctg Thr His Asn Asn Glu Leu Lys Val Leu Val Pro Val Leu Ala Val Leu 275 280 285			864
gcg gca gcg gca ttc gtt tcc ttc atc ata tac aga cgc acg cgt ccc Ala Ala Ala Ala Phe Val Ser Phe Ile Ile Tyr Arg Arg Thr Arg Pro 290 295 300			912
cac cga agc tat aca gga ccc aag act gta cag ctt gaa ctt aca gac His Arg Ser Tyr Thr Gly Pro Lys Thr Val Gln Leu Glu Leu Thr Asp 305 310 315 320			960
cac gcc His Ala			966

<210> 7
 <211> 322
 <212> PRT
 <213> mouse

<400> 7
 Met Gln Leu Lys Cys Pro Cys Phe Val Ser Leu Gly Thr Arg Gln Pro
 1 5 10 15
 Val Trp Lys Lys Leu His Val Ser Ser Gly Phe Phe Ser Gly Leu Gly
 20 25 30
 Leu Phe Leu Leu Leu Ser Ser Leu Cys Ala Ala Ser Ala Glu Thr
 35 40 45

Glu Val Gly Ala Met Val Gly Ser Asn Val Val Leu Ser Cys Ile Asp
 50 55 60
 Pro His Arg Arg His Phe Asn Leu Ser Gly Leu Tyr Val Tyr Trp Gln
 65 70 75 80
 Ile Glu Asn Pro Glu Val Ser Val Thr Tyr Tyr Leu Pro Tyr Lys Ser
 85 90 95
 Pro Gly Ile Asn Val Asp Ser Ser Tyr Lys Asn Arg Gly His Leu Ser
 100 105 110
 Leu Asp Ser Met Lys Gln Gly Asn Phe Ser Leu Tyr Leu Lys Asn Val
 115 120 125
 Thr Pro Gln Asp Thr Gln Glu Phe Thr Cys Arg Val Phe Met Asn Thr
 130 135 140
 Ala Thr Glu Leu Val Lys Ile Leu Glu Glu Val Val Arg Leu Arg Val
 145 150 155 160
 Ala Ala Asn Phe Ser Thr Pro Val Ile Ser Thr Ser Asp Ser Ser Asn
 165 170 175
 Pro Gly Gln Glu Arg Thr Tyr Thr Cys Met Ser Lys Asn Gly Tyr Pro
 180 185 190
 Glu Pro Asn Leu Tyr Trp Ile Asn Thr Thr Asp Asn Ser Leu Ile Asp
 195 200 205
 Thr Ala Leu Gln Asn Asn Thr Val Tyr Leu Asn Lys Leu Gly Leu Tyr
 210 215 220
 Asp Val Ile Ser Thr Leu Arg Leu Pro Trp Thr Ser Arg Gly Asp Val
 225 230 235 240
 Leu Cys Cys Val Glu Asn Val Ala Leu His Gln Asn Ile Thr Ser Ile
 245 250 255
 Ser Gln Ala Glu Ser Phe Thr Gly Asn Asn Thr Lys Asn Pro Gln Glu
 260 265 270
 Thr His Asn Asn Glu Leu Lys Val Leu Val Pro Val Leu Ala Val Leu
 275 280 285
 Ala Ala Ala Ala Phe Val Ser Phe Ile Ile Tyr Arg Arg Thr Arg Pro
 290 295 300
 His Arg Ser Tyr Thr Gly Pro Lys Thr Val Gln Leu Glu Leu Thr Asp
 305 310 315 320
 His Ala

<210> 8
 <211> 322
 <212> PRT
 <213> mouse

<400> 8
 Met Gln Leu Lys Cys Pro Cys Phe Val Ser Leu Gly Thr Arg Gln Pro

1	5	10	15
Val Trp Lys Lys 20	Leu His Val Ser Ser 25	Gly Phe Phe Ser Gly 30	Leu Gly
Leu Phe Leu Leu 35	Leu Ser Ser Leu 40	Cys Ala Ala Ser 45	Ala Glu Thr
Glu Val Gly Ala Met 50	Val Gly Ser Asn 55	Val Val Leu Ser 60	Cys Ile Asp
Pro His Arg Arg His 65	Phe Asn Leu Ser 70	Gly Leu Tyr Val 75	Tyr Trp Gln 80
Ile Glu Asn Pro Glu 85	Val Ser Val Thr 90	Tyr Tyr Leu Pro 95	Tyr Lys Ser 95
Pro Gly Ile Asn 100	Val Asp Ser Ser 105	Tyr Lys Asn Arg 110	Gly His Leu Ser 110
Leu Asp Ser Met Lys 115	Gln Gly Asn Phe 120	Ser Leu Tyr Leu 125	Lys Asn Val
Thr Pro Gln Asp Thr 130	Gln Glu Phe Thr 135	Cys Arg Val Phe 140	Met Asn Thr
Ala Thr Glu Leu Val 145	Lys Ile Leu Glu 150	Glu Val Val Arg 155	Leu Arg Val 160
Ala Ala Asn Phe Ser 165	Thr Pro Val Ile 170	Ser Thr Ser Asp 175	Ser Ser Asn 175
Pro Gly Gln Glu Arg 180	Thr Tyr Thr Cys 185	Met Ser Lys Asn 190	Gly Tyr Pro 190
Glu Pro Asn Leu Tyr 195	Trp Ile Asn Thr 200	Thr Asp Asn Ser 205	Leu Ile Asp 205
Thr Ala Leu Gln Asn 210	Asn Thr Val Tyr 215	Leu Asn Lys Leu 220	Gly Leu Tyr 220
Asp Val Ile Ser Thr 225	Leu Arg Leu Pro 230	Trp Thr Ser Arg 235	Gly Asp Val 240
Leu Cys Cys Val Glu 245	Asn Val Ala Leu 250	His Gln Asn Ile 255	Thr Ser Ile 255
Ser Gln Ala Glu Ser 260	Phe Thr Gly Asn 265	Asn Thr Lys Asn 270	Pro Gln Glu 270
Thr His Asn Asn Glu 275	Leu Lys Val Leu 280	Val Pro Val Leu 285	Ala Val Leu 285
Ala Ala Ala Ala Phe 290	Val Ser Phe Ile 295	Ile Tyr Arg Arg 300	Thr Arg Pro 300
His Arg Ser Tyr Thr 305	Gly Pro Lys Thr 310	Val Gln Leu Glu 315	Leu Thr Asp 320
His Ala			

<211> 306
 <212> PRT
 <213> mouse

<400> 9

```

Met Ala Cys Asn Cys Gln Leu Met Gln Asp Thr Pro Leu Leu Lys Phe
 1          5          10          15

Pro Cys Pro Arg Leu Ile Leu Leu Phe Val Leu Leu Ile Arg Leu Ser
          20          25          30

Gln Val Ser Ser Asp Val Asp Glu Gln Leu Ser Lys Ser Val Lys Asp
          35          40          45

Lys Val Leu Leu Pro Cys Arg Tyr Asn Ser Pro His Glu Asp Glu Ser
 50          55          60

Glu Asp Arg Ile Tyr Trp Gln Lys His Asp Lys Val Val Leu Ser Val
 65          70          75          80

Ile Ala Gly Lys Leu Lys Val Trp Pro Glu Tyr Lys Asn Arg Thr Leu
          85          90          95

Tyr Asp Asn Thr Thr Tyr Ser Leu Ile Ile Leu Gly Leu Val Leu Ser
          100          105          110

Asp Arg Gly Thr Tyr Ser Cys Val Val Gln Lys Lys Glu Arg Gly Thr
          115          120          125

Tyr Glu Val Lys His Leu Ala Leu Val Lys Leu Ser Ile Lys Ala Asp
          130          135          140

Phe Ser Thr Pro Asn Ile Thr Glu Ser Gly Asn Pro Ser Ala Asp Thr
          145          150          155          160

Lys Arg Ile Thr Cys Phe Ala Ser Gly Gly Phe Pro Lys Pro Arg Phe
          165          170          175

Ser Trp Leu Glu Asn Gly Arg Glu Leu Pro Gly Ile Asn Thr Thr Ile
          180          185          190

Ser Gln Asp Pro Glu Ser Glu Leu Tyr Thr Ile Ser Ser Gln Leu Asp
          195          200          205

Phe Asn Thr Thr Arg Asn His Thr Ile Lys Cys Leu Ile Lys Tyr Gly
          210          215          220

Asp Ala His Val Ser Glu Asp Phe Thr Trp Glu Lys Pro Pro Glu Asp
          225          230          235          240

Pro Pro Asp Ser Lys Asn Thr Leu Val Leu Phe Gly Ala Gly Phe Gly
          245          250          255

Ala Val Ile Thr Val Val Val Ile Val Val Ile Ile Lys Cys Phe Cys
          260          265          270

Lys His Arg Ser Cys Phe Arg Arg Asn Glu Ala Ser Arg Glu Thr Asn
          275          280          285

Asn Ser Leu Thr Phe Gly Pro Glu Glu Ala Leu Ala Glu Gln Thr Val
          290          295          300

Phe Leu
          305
  
```

<210> 10
 <211> 327
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:Synthetic
 Oglionucleotide

<400> 10
 Met Xaa Xaa Xaa Cys Xaa Cys Xaa Xaa Xaa Leu Xaa Xaa Xaa Xaa Pro
 1 5 10 15
 Xaa Xaa Xaa Xaa Leu Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Leu Xaa
 20 25 30
 Leu Phe Xaa Leu Leu Xaa Xaa Xaa Ser Xaa Xaa Xaa Xaa Xaa Xaa Xaa
 35 40 45
 Xaa Xaa Xaa Xaa Xaa Xaa Val Xaa Xaa Xaa Val Xaa Leu Xaa Cys Xaa
 50 55 60
 Xaa Xaa Xaa Xaa Xaa His Xaa Xaa Xaa Ser Xaa Xaa Xaa Xaa Tyr Trp
 65 70 75 80
 Gln Xaa Xaa Xaa Xaa Xaa Val Xaa Xaa Xaa Xaa Xaa Xaa Leu Xaa Xaa Xaa
 85 90 95
 Xaa Xaa Xaa Xaa Xaa Val Xaa Xaa Xaa Tyr Lys Asn Arg Xaa Xaa Xaa
 100 105 110
 Xaa Leu Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Ser Leu Xaa Xaa Xaa Xaa
 115 120 125
 Xaa Xaa Xaa Xaa Asp Xaa Xaa Xaa Xaa Xaa Cys Xaa Val Xaa Xaa Xaa
 130 135 140
 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Val Xaa Leu Xaa
 145 150 155 160
 Xaa Xaa Ala Xaa Phe Ser Thr Pro Xaa Ile Xaa Xaa Ser Xaa Xaa Xaa
 165 170 175
 Xaa Xaa Xaa Xaa Xaa Arg Xaa Xaa Thr Cys Xaa Xaa Xaa Xaa Gly Xaa
 180 185 190
 Pro Xaa Pro Xaa Xaa Xaa Trp Xaa Xaa Asn Xaa Xaa Xaa Xaa Xaa Xaa
 195 200 205
 Ile Xaa Thr Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
 210 215 220
 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Thr Xaa Xaa Xaa
 225 230 235 240
 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Val Xaa Xaa Xaa Xaa Xaa Xaa Xaa
 245 250 255
 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Asn Xaa Xaa Xaa Xaa Xaa
 260 265 270
 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Val Xaa Val Xaa Val Xaa Xaa

275 280 285

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Phe Xaa Xaa Xaa Xaa Xaa
 290 295 300

Xaa Arg Xaa Xaa Xaa Xaa Ser Xaa Thr Xaa Gly Pro Xaa Xaa Xaa Xaa
 305 310 315 320

Xaa Glu Xaa Thr Xaa Xaa Xaa
 325

<210> 11
 <211> 864
 <212> DNA
 <213> mouse

<220>
 <221> CDS
 <222> Complement ((1)..(864))

<400> 11

atg	cgg	ctg	ggc	agt	cct	gga	ctg	ctc	ttc	ctg	ctc	ttc	agc	agc	ctt	48
Met	Arg	Leu	Gly	Ser	Pro	Gly	Leu	Leu	Phe	Leu	Leu	Phe	Ser	Ser	Leu	
1				5					10					15		
cga	gct	gat	act	cag	gag	aag	gaa	gtc	aga	gcg	atg	gta	ggc	agc	gac	96
Arg	Ala	Asp	Thr	Gln	Glu	Lys	Glu	Val	Arg	Ala	Met	Val	Gly	Ser	Asp	
			20					25					30			
gtg	gag	ctc	agc	tgc	gct	tgc	cct	gaa	gga	agc	cgt	ttt	gat	tta	aat	144
Val	Glu	Leu	Ser	Cys	Ala	Cys	Pro	Glu	Gly	Ser	Arg	Phe	Asp	Leu	Asn	
		35					40					45				
gat	gtt	tac	gta	tat	tgg	caa	acc	agt	gag	tcg	aaa	acc	gtg	gtg	acc	192
Asp	Val	Tyr	Val	Tyr	Trp	Gln	Thr	Ser	Glu	Ser	Lys	Thr	Val	Val	Thr	
	50					55					60					
tac	cac	atc	cca	cag	aac	agc	tcc	ttg	gaa	aac	gtg	gac	agc	cgc	tac	240
Tyr	His	Ile	Pro	Gln	Asn	Ser	Ser	Leu	Glu	Asn	Val	Asp	Ser	Arg	Tyr	
	65			70						75					80	
cgg	aac	cga	gcc	ctg	atg	tca	ccg	gcc	ggc	atg	ctg	cgg	ggc	gac	ttc	288
Arg	Asn	Arg	Ala	Leu	Met	Ser	Pro	Ala	Gly	Met	Leu	Arg	Gly	Asp	Phe	
			85						90					95		
tcc	ctg	cgc	ttg	ttc	aac	gtc	acc	ccc	cag	gac	gag	cag	aag	ttt	cac	336
Ser	Leu	Arg	Leu	Phe	Asn	Val	Thr	Pro	Gln	Asp	Glu	Gln	Lys	Phe	His	
			100					105					110			
tgc	ctg	gtg	ttg	agc	caa	tcc	ctg	gga	ttc	cag	gag	gtt	ttg	agc	gtt	384
Cys	Leu	Val	Leu	Ser	Gln	Ser	Leu	Gly	Phe	Gln	Glu	Val	Leu	Ser	Val	
		115					120					125				
gag	gtt	aca	ctg	cat	gtg	gca	gca	aac	ttc	agc	gtg	ccc	gtc	gtc	agc	432
Glu	Val	Thr	Leu	His	Val	Ala	Ala	Asn	Phe	Ser	Val	Pro	Val	Val	Ser	
	130				135						140					
gcc	ccc	cac	agc	ccc	tcc	cag	gat	gag	ctc	acc	ttc	acg	tgt	aca	tcc	480
Ala	Pro	His	Ser	Pro	Ser	Gln	Asp	Glu	Leu	Thr	Phe	Thr	Cys	Thr	Ser	
	145				150					155					160	
ata	aac	ggc	tac	ccc	agg	ccc	aac	gtg	tac	tgg	atc	aat	aag	acg	gac	528
Ile	Asn	Gly	Tyr	Pro	Arg	Pro	Asn	Val	Tyr	Trp	Ile	Asn	Lys	Thr	Asp	

165										170					175					
aac	agc	ctg	ctg	gac	cag	gct	ctg	cag	aat	gac	acc	gtc	ttc	ttg	aac	576				
Asn	Ser	Leu	Leu	Asp	Gln	Ala	Leu	Gln	Asn	Asp	Thr	Val	Phe	Leu	Asn					
		180						185					190							
atg	cgg	ggc	ttg	tat	gac	gtg	gtc	agc	gtg	ctg	agg	atc	gca	cgg	acc	624				
Met	Arg	Gly	Leu	Tyr	Asp	Val	Val	Ser	Val	Leu	Arg	Ile	Ala	Arg	Thr					
		195					200					205								
ccc	agc	gtg	aac	att	ggc	tgc	tgc	ata	gag	aac	gtg	ctt	ctg	cag	cag	672				
Pro	Ser	Val	Asn	Ile	Gly	Cys	Cys	Ile	Glu	Asn	Val	Leu	Leu	Gln	Gln					
	210					215					220									
aac	ctg	act	gtc	ggc	agc	cag	aca	gga	aat	gac	atc	gga	gag	aga	gac	720				
Asn	Leu	Thr	Val	Gly	Ser	Gln	Thr	Gly	Asn	Asp	Ile	Gly	Glu	Arg	Asp					
225					230					235					240					
aag	atc	aca	gag	aat	cca	gtc	agt	acc	ggc	gag	aaa	aac	gcg	gcc	acg	768				
Lys	Ile	Thr	Glu	Asn	Pro	Val	Ser	Thr	Gly	Glu	Lys	Asn	Ala	Ala	Thr					
				245					250					255						
tgg	agc	atc	ctg	gct	gtc	ctg	tgc	ctg	ctt	gtg	gtc	gtg	gcg	gtg	gcc	816				
Trp	Ser	Ile	Leu	Ala	Val	Leu	Cys	Leu	Leu	Val	Val	Val	Ala	Val	Ala					
			260					265					270							
ata	ggc	tgg	gtg	tgc	agg	gac	cga	tgc	ctc	caa	cac	agc	tat	gca	ggt	864				
Ile	Gly	Trp	Val	Cys	Arg	Asp	Arg	Cys	Leu	Gln	His	Ser	Tyr	Ala	Gly					
		275					280					285								

<210> 12
 <211> 288
 <212> PRT
 <213> mouse

<400> 12
 Met Arg Leu Gly Ser Pro Gly Leu Leu Phe Leu Leu Phe Ser Ser Leu
 1 5 10 15
 Arg Ala Asp Thr Gln Glu Lys Glu Val Arg Ala Met Val Gly Ser Asp
 20 25 30
 Val Glu Leu Ser Cys Ala Cys Pro Glu Gly Ser Arg Phe Asp Leu Asn
 35 40 45
 Asp Val Tyr Val Tyr Trp Gln Thr Ser Glu Ser Lys Thr Val Val Thr
 50 55 60
 Tyr His Ile Pro Gln Asn Ser Ser Leu Glu Asn Val Asp Ser Arg Tyr
 65 70 75 80
 Arg Asn Arg Ala Leu Met Ser Pro Ala Gly Met Leu Arg Gly Asp Phe
 85 90 95
 Ser Leu Arg Leu Phe Asn Val Thr Pro Gln Asp Glu Gln Lys Phe His
 100 105 110
 Cys Leu Val Leu Ser Gln Ser Leu Gly Phe Gln Glu Val Leu Ser Val
 115 120 125
 Glu Val Thr Leu His Val Ala Ala Asn Phe Ser Val Pro Val Val Ser
 130 135 140

Ala Pro His Ser Pro Ser Gln Asp Glu Leu Thr Phe Thr Cys Thr Ser
 145 150 155 160
 Ile Asn Gly Tyr Pro Arg Pro Asn Val Tyr Trp Ile Asn Lys Thr Asp
 165 170 175
 Asn Ser Leu Leu Asp Gln Ala Leu Gln Asn Asp Thr Val Phe Leu Asn
 180 185 190
 Met Arg Gly Leu Tyr Asp Val Val Ser Val Leu Arg Ile Ala Arg Thr
 195 200 205
 Pro Ser Val Asn Ile Gly Cys Cys Ile Glu Asn Val Leu Leu Gln Gln
 210 215 220
 Asn Leu Thr Val Gly Ser Gln Thr Gly Asn Asp Ile Gly Glu Arg Asp
 225 230 235 240
 Lys Ile Thr Glu Asn Pro Val Ser Thr Gly Glu Lys Asn Ala Ala Thr
 245 250 255
 Trp Ser Ile Leu Ala Val Leu Cys Leu Leu Val Val Val Ala Val Ala
 260 265 270
 Ile Gly Trp Val Cys Arg Asp Arg Cys Leu Gln His Ser Tyr Ala Gly
 275 280 285

<210> 13
 <211> 267
 <212> PRT
 <213> Human

<400> 13
 Glu Lys Glu Val Arg Ala Met Val Gly Ser Asp Val Glu Leu Ser Cys
 1 5 10 15
 Ala Cys Pro Glu Gly Ser Arg Phe Asp Leu Asn Asp Val Tyr Val Tyr
 20 25 30
 Trp Gln Thr Ser Glu Ser Lys Thr Val Val Thr Tyr His Ile Pro Gln
 35 40 45
 Asn Ser Ser Leu Glu Asn Val Asp Ser Arg Tyr Arg Asn Arg Ala Leu
 50 55 60
 Met Ser Pro Ala Gly Met Leu Arg Gly Asp Phe Ser Leu Arg Leu Phe
 65 70 75 80
 Asn Val Thr Pro Gln Asp Glu Gln Lys Phe His Cys Leu Val Leu Ser
 85 90 95
 Gln Ser Leu Gly Phe Gln Glu Val Leu Ser Val Glu Val Thr Leu His
 100 105 110
 Val Ala Ala Asn Phe Ser Val Pro Val Val Ser Ala Pro His Ser Pro
 115 120 125
 Ser Gln Asp Glu Leu Thr Phe Thr Cys Thr Ser Ile Asn Gly Tyr Pro
 130 135 140
 Arg Pro Asn Val Tyr Trp Ile Asn Lys Thr Asp Asn Ser Leu Leu Asp
 145 150 155 160

Gln	Ala	Leu	Gln	Asn 165	Asp	Thr	Val	Phe	Leu 170	Asn	Met	Arg	Gly	Leu 175	Tyr
Asp	Val	Val	Ser 180	Val	Leu	Arg	Ile	Ala 185	Arg	Thr	Pro	Ser	Val 190	Asn	Ile
Gly	Cys	Cys 195	Ile	Glu	Asn	Val	Leu 200	Leu	Gln	Gln	Asn	Leu 205	Thr	Val	Gly
Ser	Gln 210	Thr	Gly	Asn	Asp	Ile 215	Gly	Glu	Arg	Asp	Lys 220	Ile	Thr	Glu	Asn
Pro 225	Val	Ser	Thr	Gly	Glu 230	Lys	Asn	Ala	Ala	Thr 235	Trp	Ser	Ile	Leu	Ala 240
Val	Leu	Cys	Leu	Leu 245	Val	Val	Val	Ala	Val 250	Ala	Ile	Gly	Trp	Val 255	Cys
Arg	Asp	Arg	Cys 260	Leu	Gln	His	Ser	Tyr 265	Ala	Gly					

```
<210> 14
<211> 276
<212> PRT
<213> mouse
```

<400>	14															
Glu	Thr	Glu	Val	Gly	Ala	Met	Val	Gly	Ser	Asn	Val	Val	Leu	Ser	Cys	
1				5					10					15		
Ile	Asp	Pro	His	Arg	Arg	His	Phe	Asn	Leu	Ser	Gly	Leu	Tyr	Val	Tyr	
			20					25					30			
Trp	Gln	Ile	Glu	Asn	Pro	Glu	Val	Ser	Val	Thr	Tyr	Tyr	Leu	Pro	Tyr	
		35					40					45				
Lys	Ser	Pro	Gly	Ile	Asn	Val	Asp	Ser	Ser	Tyr	Lys	Asn	Arg	Gly	His	
50						55					60					
Leu	Ser	Leu	Asp	Ser	Met	Lys	Gln	Gly	Asn	Phe	Ser	Leu	Tyr	Leu	Lys	
65					70					75					80	
Asn	Val	Thr	Pro	Gln	Asp	Thr	Gln	Glu	Phe	Thr	Cys	Arg	Val	Phe	Met	
				85					90					95		
Asn	Thr	Ala	Thr	Glu	Leu	Val	Lys	Ile	Leu	Glu	Glu	Val	Val	Arg	Leu	
			100					105					110			
Arg	Val	Ala	Ala	Asn	Phe	Ser	Thr	Pro	Val	Ile	Ser	Thr	Ser	Asp	Ser	
		115					120					125				
Ser	Asn	Pro	Gly	Gln	Glu	Arg	Thr	Tyr	Thr	Cys	Met	Ser	Lys	Asn	Gly	
130						135					140					
Tyr	Pro	Glu	Pro	Asn	Leu	Tyr	Trp	Ile	Asn	Thr	Thr	Asp	Asn	Ser	Leu	
145					150					155					160	
Ile	Asp	Thr	Ala	Leu	Gln	Asn	Asn	Thr	Val	Tyr	Leu	Asn	Lys	Leu	Gly	
				165					170					175		
Leu	Tyr	Asp	Val	Ile	Ser	Thr	Leu	Arg	Leu	Pro	Trp	Thr	Ser	Arg	Gly	
			180					185					190			

Asp Val Leu Cys Cys Val Glu Asn Val Ala Leu His Gln Asn Ile Thr
 195 200 205
 Ser Ile Ser Gln Ala Glu Ser Phe Thr Gly Asn Asn Thr Lys Asn Pro
 210 215 220
 Gln Glu Thr His Asn Asn Glu Leu Lys Val Leu Val Pro Val Leu Ala
 225 230 235 240
 Val Leu Ala Ala Ala Ala Phe Val Ser Phe Ile Ile Tyr Arg Arg Thr
 245 250 255
 Arg Pro His Arg Ser Tyr Thr Gly Pro Lys Thr Val Gln Leu Glu Leu
 260 265 270
 Thr Asp His Ala
 275

<210> 15
 <211> 280
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:Synthetic
 Oglionucleotide

<400> 15
 Glu Xaa Glu Val Xaa Ala Met Gly Val Ser Xaa Val Xaa Leu Ser Cys
 1 5 10 15
 Xaa Xaa Pro Xaa Xaa Xaa Xaa Phe Xaa Leu Xaa Xaa Xaa Tyr Val Tyr
 20 25 30
 Trp Gln Xaa Xaa Xaa Xaa Xaa Xaa Xaa Val Thr Tyr Xaa Xaa Pro Xaa
 35 40 45
 Xaa Ser Xaa Xaa Xaa Asn Val Asp Ser Xaa Tyr Xaa Asn Arg Xaa Xaa
 50 55 60
 Xaa Ser Xaa Xaa Xaa Met Xaa Xaa Gly Xaa Phe Ser Leu Xaa Leu Xaa
 65 70 75 80
 Asn Val Thr Pro Gln Asp Xaa Gln Xaa Phe Xaa Cys Xaa Val Xaa Xaa
 85 90 95
 Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Leu Xaa Xaa Xaa Val Xaa Leu
 100 105 110
 Xaa Val Ala Ala Asn Phe Ser Xaa Pro Val Xaa Ser Xaa Xaa Xaa Ser
 115 120 125
 Xaa Xaa Xaa Xaa Xaa Glu Xaa Thr Xaa Thr Cys Xaa Ser Xaa Asn Gly
 130 135 140
 Tyr Pro Xaa Pro Asn Xaa Tyr Trp Ile Asn Xaa Thr Asp Asn Ser Leu
 145 150 155 160
 Xaa Asp Xaa Ala Leu Gln Asn Xaa Thr Val Xaa Leu Asn Xaa Xaa Gly
 165 170 175
 Leu Tyr Asp Val Xaa Ser Xaa Leu Arg Xaa Xaa Xaa Thr Xaa Xaa Xaa

	180		185		190
Xaa Xaa Xaa Cys Cys Xaa Glu Asn Val Xaa Leu Xaa Gln Asn Xaa Thr					
195		200		205	
Xaa Xaa Ser Gln Xaa Xaa Xaa Xaa Xaa Gly Xaa Xaa Lys Xaa Xaa Xaa					
210		215		220	
Xaa Xaa Xaa Xaa Xaa Xaa Xaa Lys Xaa Xaa Xaa Xaa Xaa Xaa Xaa Leu					
225		230		235	240
Ala Val Leu Xaa Xaa Xaa Xaa Xaa Val Xaa Xaa Xaa Ile Xaa Xaa Xaa					
	245		250		255
Xaa Arg Xaa Arg Xaa Xaa Xaa Xaa Ser Tyr Xaa Gly Xaa Xaa Xaa Xaa					
	260		265		270
Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa					
	275		280		

<210> 16
 <211> 1294
 <212> DNA
 <213> Human

<220>
 <221> 5'UTR
 <222> (1)..(199)

<220>
 <221> CDS
 <222> (200)..(1105)

<400> 16
 gctggtacgc ctgcaggtac cgggtccggaa ttcccgggtc gaccacgcgc tccgcccacg 60
 cgctccgcggg agcgcagtta gagccgatct cccgcgcccc gaggttgctc ctctccgagg 120
 tctcccgcg cccaagttct ccgcgccccg aggtctccgc gccccgaggt ctccgcggcc 180
 cgaggtctcc gcccgcacc atg cgg ctg ggc agt cct gga ctg ctc ttc ctg 232
 Met Arg Leu Gly Ser Pro Gly Leu Leu Phe Leu
 1 5 10
 ctc ttc agc agc ctt cga gct gat act cag gag aag gaa gtc aga gcg 280
 Leu Phe Ser Ser Leu Arg Ala Asp Thr Gln Glu Lys Glu Val Arg Ala
 15 20 25
 atg gta ggc agc gac gtg gag ctc agc tgc gct tgc cct gaa gga agc 328
 Met Val Gly Ser Asp Val Glu Leu Ser Cys Ala Cys Pro Glu Gly Ser
 30 35 40
 cgt ttt gat tta aat gat gtt tac gta tat tgg caa acc agt gag tcg 376
 Arg Phe Asp Leu Asn Asp Val Tyr Val Tyr Trp Gln Thr Ser Glu Ser
 45 50 55
 aaa acc gtg gtg acc tac cac atc cca cag aac agc tcc ttg gaa aac 424
 Lys Thr Val Val Thr Tyr His Ile Pro Gln Asn Ser Ser Leu Glu Asn
 60 65 70 75
 gtg gac agc cgc tac cgg aac cga gcc ctg atg tca ccg gcc ggc atg 472
 Val Asp Ser Arg Tyr Arg Asn Arg Ala Leu Met Ser Pro Ala Gly Met
 80 85 90

ctg cgg ggc gac ttc tcc ctg cgc ttg ttc aac gtc acc ccc cag gac 520
 Leu Arg Gly Asp Phe Ser Leu Arg Leu Phe Asn Val Thr Pro Gln Asp
 95 100 105

gag cag aag ttt cac tgc ctg gtg ttg agc caa tcc ctg gga ttc cag 568
 Glu Gln Lys Phe His Cys Leu Val Leu Ser Gln Ser Leu Gly Phe Gln
 110 115 120

gag gtt ttg agc gtt gag gtt aca ctg cat gtg gca gca aac ttc agc 616
 Glu Val Leu Ser Val Glu Val Thr Leu His Val Ala Ala Asn Phe Ser
 125 130 135

gtg ccc gtc gtc agc gcc ccc cac agc ccc tcc cag gat gag ctc acc 664
 Val Pro Val Val Ser Ala Pro His Ser Pro Ser Gln Asp Glu Leu Thr
 140 145 150 155

ttc acg tgt aca tcc ata aac ggc tac ccc agg ccc aac gtg tac tgg 712
 Phe Thr Cys Thr Ser Ile Asn Gly Tyr Pro Arg Pro Asn Val Tyr Trp
 160 165 170

atc aat aag acg gac aac agc ctg ctg gac cag gct ctg cag aat gac 760
 Ile Asn Lys Thr Asp Asn Ser Leu Leu Asp Gln Ala Leu Gln Asn Asp
 175 180 185

acc gtc ttc ttg aac atg cgg ggc ttg tat gac gtg gtc agc gtg ctg 808
 Thr Val Phe Leu Asn Met Arg Gly Leu Tyr Asp Val Val Ser Val Leu
 190 195 200

agg atc gca cgg acc ccc agc gtg aac att ggc tgc tgc ata gag aac 856
 Arg Ile Ala Arg Thr Pro Ser Val Asn Ile Gly Cys Cys Ile Glu Asn
 205 210 215

gtg ctt ctg cag cag aac ctg act gtc ggc agc cag aca gga aat gac 904
 Val Leu Leu Gln Gln Asn Leu Thr Val Gly Ser Gln Thr Gly Asn Asp
 220 225 230 235

atc gga gag aga gac aag atc aca gag aat cca gtc agt acc ggc gag 952
 Ile Gly Glu Arg Asp Lys Ile Thr Glu Asn Pro Val Ser Thr Gly Glu
 240 245 250

aaa aac gcg gcc acg tgg agc atc ctg gct gtc ctg tgc ctg ctt gtg 1000
 Lys Asn Ala Ala Thr Trp Ser Ile Leu Ala Val Leu Cys Leu Leu Val
 255 260 265

gtc gtg gcg gtg gcc ata ggc tgg gtg tgc agg gac cga tgc ctc caa 1048
 Val Val Ala Val Ala Ile Gly Trp Val Cys Arg Asp Arg Cys Leu Gln
 270 275 280

cac agc tat gca ggt gcc tgg gct gtg agt ccg gag aca gag ctc act 1096
 His Ser Tyr Ala Gly Ala Trp Ala Val Ser Pro Glu Thr Glu Leu Thr
 285 290 295

ggc cac gtt tgaccggagc tcaccgccca gagcgtggac agggcttccg 1145
 Gly His Val
 300

tgagacgccca ccgtgagagg ccaggtggca gcttgagcat ggactcccag actgcagggg 1205

agcacttggg gcagccccca gaaggaccac tgctggatcc cagggagaac ctgctggcgt 1265

tggctgtgat cctggaatga ggccctttc 1294

<210> 17
 <211> 302
 <212> PRT
 <213> Human

<400> 17

```

Met Arg Leu Gly Ser Pro Gly Leu Leu Phe Leu Leu Phe Ser Ser Leu
 1          5          10          15

Arg Ala Asp Thr Gln Glu Lys Glu Val Arg Ala Met Val Gly Ser Asp
      20          25          30

Val Glu Leu Ser Cys Ala Cys Pro Glu Gly Ser Arg Phe Asp Leu Asn
      35          40          45

Asp Val Tyr Val Tyr Trp Gln Thr Ser Glu Ser Lys Thr Val Val Thr
      50          55          60

Tyr His Ile Pro Gln Asn Ser Ser Leu Glu Asn Val Asp Ser Arg Tyr
      65          70          75          80

Arg Asn Arg Ala Leu Met Ser Pro Ala Gly Met Leu Arg Gly Asp Phe
      85          90          95

Ser Leu Arg Leu Phe Asn Val Thr Pro Gln Asp Glu Gln Lys Phe His
      100          105          110

Cys Leu Val Leu Ser Gln Ser Leu Gly Phe Gln Glu Val Leu Ser Val
      115          120          125

Glu Val Thr Leu His Val Ala Ala Asn Phe Ser Val Pro Val Val Ser
      130          135          140

Ala Pro His Ser Pro Ser Gln Asp Glu Leu Thr Phe Thr Cys Thr Ser
      145          150          155          160

Ile Asn Gly Tyr Pro Arg Pro Asn Val Tyr Trp Ile Asn Lys Thr Asp
      165          170          175

Asn Ser Leu Leu Asp Gln Ala Leu Gln Asn Asp Thr Val Phe Leu Asn
      180          185          190

Met Arg Gly Leu Tyr Asp Val Val Ser Val Leu Arg Ile Ala Arg Thr
      195          200          205

Pro Ser Val Asn Ile Gly Cys Cys Ile Glu Asn Val Leu Leu Gln Gln
      210          215          220

Asn Leu Thr Val Gly Ser Gln Thr Gly Asn Asp Ile Gly Glu Arg Asp
      225          230          235          240

Lys Ile Thr Glu Asn Pro Val Ser Thr Gly Glu Lys Asn Ala Ala Thr
      245          250          255

Trp Ser Ile Leu Ala Val Leu Cys Leu Leu Val Val Val Ala Val Ala
      260          265          270

Ile Gly Trp Val Cys Arg Asp Arg Cys Leu Gln His Ser Tyr Ala Gly
      275          280          285

Ala Trp Ala Val Ser Pro Glu Thr Glu Leu Thr Gly His Val
      290          295          300

```

<210> 18
 <211> 302
 <212> PRT
 <213> Human

<400> 18

```

Met Arg Leu Gly Ser Pro Gly Leu Leu Phe Leu Leu Phe Ser Ser Leu
 1           5           10           15

Arg Ala Asp Thr Gln Glu Lys Glu Val Arg Ala Met Val Gly Ser Asp
          20           25           30

Val Glu Leu Ser Cys Ala Cys Pro Glu Gly Ser Arg Phe Asp Leu Asn
          35           40           45

Asp Val Tyr Val Tyr Trp Gln Thr Ser Glu Ser Lys Thr Val Val Thr
 50           55           60

Tyr His Ile Pro Gln Asn Ser Ser Leu Glu Asn Val Asp Ser Arg Tyr
 65           70           75           80

Arg Asn Arg Ala Leu Met Ser Pro Ala Gly Met Leu Arg Gly Asp Phe
          85           90           95

Ser Leu Arg Leu Phe Asn Val Thr Pro Gln Asp Glu Gln Lys Phe His
          100          105          110

Cys Leu Val Leu Ser Gln Ser Leu Gly Phe Gln Glu Val Leu Ser Val
          115          120          125

Glu Val Thr Leu His Val Ala Ala Asn Phe Ser Val Pro Val Val Ser
          130          135          140

Ala Pro His Ser Pro Ser Gln Asp Glu Leu Thr Phe Thr Cys Thr Ser
          145          150          155          160

Ile Asn Gly Tyr Pro Arg Pro Asn Val Tyr Trp Ile Asn Lys Thr Asp
          165          170          175

Asn Ser Leu Leu Asp Gln Ala Leu Gln Asn Asp Thr Val Phe Leu Asn
          180          185          190

Met Arg Gly Leu Tyr Asp Val Val Ser Val Leu Arg Ile Ala Arg Thr
          195          200          205

Pro Ser Val Asn Ile Gly Cys Cys Ile Glu Asn Val Leu Leu Gln Gln
          210          215          220

Asn Leu Thr Val Gly Ser Gln Thr Gly Asn Asp Ile Gly Glu Arg Asp
          225          230          235          240

Lys Ile Thr Glu Asn Pro Val Ser Thr Gly Glu Lys Asn Ala Ala Thr
          245          250          255

Trp Ser Ile Leu Ala Val Leu Cys Leu Leu Val Val Val Ala Val Ala
          260          265          270

Ile Gly Trp Val Cys Arg Asp Arg Cys Leu Gln His Ser Tyr Ala Gly
          275          280          285

Ala Trp Ala Val Ser Pro Glu Thr Glu Leu Thr Gly His Val
          290          295          300

```

```
<210> 19
<211> 322
<212> PRT
<213> mouse
```

<400> 19

Met 1	Gln	Leu	Lys	Cys 5	Pro	Cys	Phe	Val	Ser 10	Leu	Gly	Thr	Arg	Gln	Pro 15
Val	Trp	Lys	Lys 20	Leu	His	Val	Ser	Ser 25	Gly	Phe	Phe	Ser	Gly 30	Leu	Gly
Leu	Phe	Leu 35	Leu	Leu	Leu	Ser	Ser 40	Leu	Cys	Ala	Ala	Ser 45	Ala	Glu	Thr
Glu	Val 50	Gly	Ala	Met	Val	Gly 55	Ser	Asn	Val	Val	Leu 60	Ser	Cys	Ile	Asp
Pro 65	His	Arg	Arg	His	Phe 70	Asn	Leu	Ser	Gly	Leu 75	Tyr	Val	Tyr	Trp	Gln 80
Ile	Glu	Asn	Pro	Glu 85	Val	Ser	Val	Thr	Tyr 90	Tyr	Leu	Pro	Tyr	Lys 95	Ser
Pro	Gly	Ile	Asn 100	Val	Asp	Ser	Ser	Tyr 105	Lys	Asn	Arg	Gly	His 110	Leu	Ser
Leu	Asp	Ser 115	Met	Lys	Gln	Gly	Asn 120	Phe	Ser	Leu	Tyr	Leu 125	Lys	Asn	Val
Thr	Pro 130	Gln	Asp	Thr	Gln	Glu 135	Phe	Thr	Cys	Arg	Val 140	Phe	Met	Asn	Thr
Ala 145	Thr	Glu	Leu	Val	Lys 150	Ile	Leu	Glu	Glu	Val 155	Val	Arg	Leu	Arg	Val 160
Ala	Ala	Asn	Phe	Ser 165	Thr	Pro	Val	Ile	Ser 170	Thr	Ser	Asp	Ser	Ser 175	Asn
Pro	Gly	Gln	Glu 180	Arg	Thr	Tyr	Thr	Cys 185	Met	Ser	Lys	Asn	Gly 190	Tyr	Pro
Glu	Pro	Asn 195	Leu	Tyr	Trp	Ile	Asn 200	Thr	Thr	Asp	Asn	Ser 205	Leu	Ile	Asp
Thr	Ala 210	Leu	Gln	Asn	Asn	Thr 215	Val	Tyr	Leu	Asn	Lys 220	Leu	Gly	Leu	Tyr
Asp 225	Val	Ile	Ser	Thr	Leu 230	Arg	Leu	Pro	Trp	Thr 235	Ser	Arg	Gly	Asp	Val 240
Leu	Cys	Cys	Val	Glu 245	Asn	Val	Ala	Leu	His 250	Gln	Asn	Ile	Thr	Ser 255	Ile
Ser	Gln	Ala	Glu 260	Ser	Phe	Thr	Gly	Asn 265	Asn	Thr	Lys	Asn	Pro 270	Gln	Glu
Thr	His 275	Asn	Asn	Glu	Leu	Lys	Val 280	Leu	Val	Pro	Val	Leu 285	Ala	Val	Leu
Ala	Ala 290	Ala	Ala	Phe	Val	Ser 295	Phe	Ile	Ile	Tyr	Arg 300	Arg	Thr	Arg	Pro

His Arg Ser Tyr Thr Gly Pro Lys Thr Val Gln Leu Glu Leu Thr Asp
 305 310 315 320

His Ala

<210> 20

<211> 329

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic
 Oglionucleotide

<400> 20

Met Xaa Leu Xaa Xaa Pro Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
 1 5 10 15

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Gly Leu Xaa
 20 25 30

Leu Phe Xaa Leu Leu Xaa Ser Ser Leu Xaa Ala Xaa Xaa Xaa Glu Xaa
 35 40 45

Glu Val Xaa Ala Met Val Gly Ser Xaa Val Xaa Leu Ser Cys Xaa Xaa
 50 55 60

Pro Xaa Xaa Xaa Xaa Phe Xaa Leu Xaa Xaa Xaa Tyr Val Tyr Trp Gln
 65 70 75 80

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Val Thr Tyr Xaa Xaa Pro Xaa Xaa Ser
 85 90 95

Xaa Xaa Xaa Asn Val Asp Ser Xaa Tyr Xaa Asn Arg Xaa Xaa Xaa Ser
 100 105 110

Xaa Xaa Xaa Met Xaa Xaa Gly Xaa Phe Ser Leu Xaa Leu Xaa Asn Val
 115 120 125

Thr Pro Gln Asp Xaa Gln Xaa Phe Xaa Cys Xaa Val Xaa Xaa Xaa Xaa
 130 135 140

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Leu Xaa Xaa Xaa Val Xaa Leu Xaa Val
 145 150 155 160

Ala Ala Asn Phe Ser Xaa Pro Val Xaa Ser Xaa Xaa Xaa Ser Xaa Xaa
 165 170 175

Xaa Xaa Xaa Glu Xaa Thr Xaa Thr Cys Xaa Ser Xaa Asn Gly Tyr Pro
 180 185 190

Xaa Pro Asn Xaa Tyr Trp Ile Asn Xaa Thr Asp Asn Ser Leu Xaa Asp
 195 200 205

Xaa Ala Leu Gln Asn Xaa Thr Val Xaa Leu Asn Xaa Xaa Gly Leu Tyr
 210 215 220

Asp Val Xaa Ser Xaa Leu Arg Xaa Xaa Xaa Thr Xaa Xaa Xaa Xaa Xaa
 225 230 235 240

Xaa Cys Cys Xaa Glu Asn Val Xaa Leu Xaa Gln Asn Xaa Thr Xaa Xaa
 245 250 255

Ser Gln Xaa Xaa Xaa Xaa Xaa Gly Xaa Xaa Xaa Lys Xaa Xaa Xaa Xaa
 260 265 270
 Xaa Xaa Xaa Xaa Xaa Xaa Lys Xaa Xaa Xaa Xaa Xaa Xaa Xaa Leu Ala
 275 280 285
 Val Leu Xaa Xaa Xaa Xaa Xaa Val Xaa Xaa Xaa Ile Xaa Xaa Xaa Xaa
 290 295 300
 Arg Xaa Arg Xaa Xaa Xaa Xaa Ser Tyr Xaa Gly Xaa Xaa Xaa Val Xaa
 305 310 315 320
 Xaa Glu Xaa Xaa Leu Thr Xaa His Xaa
 325

<210> 21
 <211> 1370
 <212> DNA
 <213> Human

<220>
 <221> 5'UTR
 <222> (1)..(165)

<220>
 <221> CDS
 <222> (166)..(762)

<400> 21
 aacaatttca cacaggaaac agctatgacc atgattacgc caagctctaa tacgactcac 60
 tatagggaaa gctgggtacgc ctgcaggtac cgggtccggaa ttcccgggtc gaccacgcgc 120
 tccgtgaaca ctgaacgcga ggactgttaa ctgtttctgg caaac atg aag tca ggc 177
 Met Lys Ser Gly
 1
 ctc tgg tat ttc ttt ctc ttc tgc ttg cgc att aaa gtt tta aca gga 225
 Leu Trp Tyr Phe Phe Leu Phe Cys Leu Arg Ile Lys Val Leu Thr Gly
 5 10 15 20
 gaa atc aat ggt tct gcc aat tat gag atg ttt ata ttt cac aac gga 273
 Glu Ile Asn Gly Ser Ala Asn Tyr Glu Met Phe Ile Phe His Asn Gly
 25 30 35
 ggt gta caa att tta tgc aaa tat cct gac att gtc cag caa ttt aaa 321
 Gly Val Gln Ile Leu Cys Lys Tyr Pro Asp Ile Val Gln Gln Phe Lys
 40 45 50
 atg cag ttg ctg aaa ggg ggg caa ata ctc tgc gat ctc act aag aca 369
 Met Gln Leu Leu Lys Gly Gly Gln Ile Leu Cys Asp Leu Thr Lys Thr
 55 60 65
 aaa gga agt gga aac aca gtg tcc att aag agt ctg aaa ttc tgc cat 417
 Lys Gly Ser Gly Asn Thr Val Ser Ile Lys Ser Leu Lys Phe Cys His
 70 75 80
 tct cag tta tcc aac aac agt gtc tct ttt ttt cta tac aac ttg gac 465
 Ser Gln Leu Ser Asn Asn Ser Val Ser Phe Phe Leu Tyr Asn Leu Asp
 85 90 95 100
 cat tct cat gcc aac tat tac ttc tgc aac cta tca att ttt gat cct 513

His Ser His Ala Asn Tyr Tyr Phe Cys Asn Leu Ser Ile Phe Asp Pro
 105 110 115
 cct cct ttt aaa gta act ctt aca gga gga tat ttg cat att tat gaa 561
 Pro Pro Phe Lys Val Thr Leu Thr Gly Gly Tyr Leu His Ile Tyr Glu
 120 125 130
 tca caa ctt tgt tgc cag ctg aag ttc tgg tta ccc ata gga tgt gca 609
 Ser Gln Leu Cys Cys Gln Leu Lys Phe Trp Leu Pro Ile Gly Cys Ala
 135 140 145
 gcc ttt gtt gta gtc tgc att ttg gga tgc ata ctt att tgt tgg ctt 657
 Ala Phe Val Val Val Cys Ile Leu Gly Cys Ile Leu Ile Cys Trp Leu
 150 155 160
 aca aaa aag aag tat tca tcc agt gtg cac gac cct aac ggt gaa tac 705
 Thr Lys Lys Lys Tyr Ser Ser Val His Asp Pro Asn Gly Glu Tyr
 165 170 175 180
 atg ttc atg aga gca gtg aac aca gcc aaa aaa tct aga ctc aca gat 753
 Met Phe Met Arg Ala Val Asn Thr Ala Lys Lys Ser Arg Leu Thr Asp
 185 190 195
 gtg acc cta taatatggaa ctctggcacc caggcatgaa gcacgttggc 802
 Val Thr Leu
 cagttttcct caacttgaag tgcaagattc tcttatttcc gggaccacgg agagtctgac 862
 ttaactacat acatcttctg ctggtgtttt gttcaatctg gaagaatgac tgtatcagtc 922
 aatggggatt ttaacagact gccttggtac tgccgagtc tctcaaaaca aacaccctct 982
 tgcaaccagc tttggagaaa gccagctcc tgtgtgctca ctgggagtg aatccctgtc 1042
 tccacatctg ctctagcag tgcacagcc agtaaaacaa acacatttac aagaaaaatg 1102
 ttttaaagat gccaggggta ctgaatctgc aaagcaaag agcagccaag gaccagcatc 1162
 tgtccgcatt tcactatcat actacctctt ctttctgtag ggatgagaat tcctctttta 1222
 atcagtcaag ggagatgctt caaagctgga gctattttat ttctgagatg ttgatgtgaa 1282
 ctgtacatta gtacatactc agtactctcc ttcaattgct gaaccccagt tgaccatttt 1342
 accaagactt tagatgcttt cttgtgcc 1370

 <210> 22
 <211> 199
 <212> PRT
 <213> Human

 <400> 22
 Met Lys Ser Gly Leu Trp Tyr Phe Phe Leu Phe Cys Leu Arg Ile Lys
 1 5 10 15
 Val Leu Thr Gly Glu Ile Asn Gly Ser Ala Asn Tyr Glu Met Phe Ile
 20 25 30
 Phe His Asn Gly Gly Val Gln Ile Leu Cys Lys Tyr Pro Asp Ile Val
 35 40 45
 Gln Gln Phe Lys Met Gln Leu Leu Lys Gly Gly Gln Ile Leu Cys Asp
 50 55 60

Leu Thr Lys Thr Lys Gly Ser Gly Asn Thr Val Ser Ile Lys Ser Leu
 65 70 75 80
 Lys Phe Cys His Ser Gln Leu Ser Asn Asn Ser Val Ser Phe Phe Leu
 85 90 95
 Tyr Asn Leu Asp His Ser His Ala Asn Tyr Tyr Phe Cys Asn Leu Ser
 100 105 110
 Ile Phe Asp Pro Pro Pro Phe Lys Val Thr Leu Thr Gly Gly Tyr Leu
 115 120 125
 His Ile Tyr Glu Ser Gln Leu Cys Cys Gln Leu Lys Phe Trp Leu Pro
 130 135 140
 Ile Gly Cys Ala Ala Phe Val Val Val Cys Ile Leu Gly Cys Ile Leu
 145 150 155 160
 Ile Cys Trp Leu Thr Lys Lys Lys Tyr Ser Ser Ser Val His Asp Pro
 165 170 175
 Asn Gly Glu Tyr Met Phe Met Arg Ala Val Asn Thr Ala Lys Lys Ser
 180 185 190
 Arg Leu Thr Asp Val Thr Leu
 195

<210> 23
 <211> 199
 <212> PRT
 <213> Human

<400> 23
 Met Lys Ser Gly Leu Trp Tyr Phe Phe Leu Phe Cys Leu Arg Ile Lys
 1 5 10 15
 Val Leu Thr Gly Glu Ile Asn Gly Ser Ala Asn Tyr Glu Met Phe Ile
 20 25 30
 Phe His Asn Gly Gly Val Gln Ile Leu Cys Lys Tyr Pro Asp Ile Val
 35 40 45
 Gln Gln Phe Lys Met Gln Leu Leu Lys Gly Gly Gln Ile Leu Cys Asp
 50 55 60
 Leu Thr Lys Thr Lys Gly Ser Gly Asn Thr Val Ser Ile Lys Ser Leu
 65 70 75 80
 Lys Phe Cys His Ser Gln Leu Ser Asn Asn Ser Val Ser Phe Phe Leu
 85 90 95
 Tyr Asn Leu Asp His Ser His Ala Asn Tyr Tyr Phe Cys Asn Leu Ser
 100 105 110
 Ile Phe Asp Pro Pro Pro Phe Lys Val Thr Leu Thr Gly Gly Tyr Leu
 115 120 125
 His Ile Tyr Glu Ser Gln Leu Cys Cys Gln Leu Lys Phe Trp Leu Pro
 130 135 140

Ile Gly Cys Ala Ala Phe Val Val Val Cys Ile Leu Gly Cys Ile Leu
 145 150 155 160
 Ile Cys Trp Leu Thr Lys Lys Lys Tyr Ser Ser Ser Val His Asp Pro
 165 170 175
 Asn Gly Glu Tyr Met Phe Met Arg Ala Val Asn Thr Ala Lys Lys Ser
 180 185 190
 Arg Leu Thr Asp Val Thr Leu
 195

<210> 24
 <211> 200
 <212> PRT
 <213> mouse

<400> 24
 Met Lys Pro Tyr Phe Cys Arg Val Phe Val Phe Cys Phe Leu Ile Arg
 1 5 10 15
 Leu Leu Thr Gly Glu Ile Asn Gly Ser Ala Asp His Arg Met Phe Ser
 20 25 30
 Phe His Asn Gly Gly Val Gln Ile Ser Cys Lys Tyr Pro Glu Thr Val
 35 40 45
 Gln Gln Leu Lys Met Arg Leu Phe Arg Glu Arg Glu Val Leu Cys Glu
 50 55 60
 Leu Thr Lys Thr Lys Gly Ser Gly Asn Ala Val Ser Ile Lys Asn Pro
 65 70 75 80
 Met Leu Cys Leu Tyr His Leu Ser Asn Asn Ser Val Ser Phe Phe Leu
 85 90 95
 Asn Asn Pro Asp Ser Ser Gln Gly Ser Tyr Tyr Phe Cys Ser Leu Ser
 100 105 110
 Ile Phe Asp Pro Pro Pro Phe Gln Glu Arg Asn Leu Ser Gly Gly Tyr
 115 120 125
 Leu His Ile Tyr Glu Ser Gln Leu Cys Cys Gln Leu Lys Leu Trp Leu
 130 135 140
 Pro Val Gly Cys Ala Ala Phe Val Val Val Leu Leu Phe Gly Cys Ile
 145 150 155 160
 Leu Ile Ile Trp Phe Ser Lys Lys Lys Tyr Gly Ser Ser Val His Asp
 165 170 175
 Pro Asn Ser Glu Tyr Met Phe Met Ala Ala Val Asn Thr Asn Lys Lys
 180 185 190
 Ser Arg Leu Ala Gly Val Thr Ser
 195 200

<210> 25
 <211> 24
 <212> DNA
 <213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:Synthetic
Oglionucleotide

<400> 25
accatgcggc tgggcagtcg tgga 24

<210> 26
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:Synthetic
Oglionucleotide

<400> 26
tggtgaccta ccacatccca cag 23

<210> 27
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:Synthetic
Oglionucleotide

<400> 27
tccgatgtca tttcctgtct ggc 23

<210> 28
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:Synthetic
Oglionucleotide

<400> 28
gctctgtctc cggactcaca gccc 24

<210> 29
<211> 28
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:Synthetic
Oglionucleotide

<400> 29
gtggcagcaa acttcagcgt gcccgtcg 28

<210> 30
<211> 28
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:Synthetic
Oglionucleotide

<400> 30
cccaacgtgt actggatcaa taagacgg 28

<210> 31
<211> 28
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:Synthetic
Oglionucleotide

<400> 31
gcgtgctgag gatcgacgg acccccag 28

<210> 32
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:Synthetic
Oglionucleotide

<400> 32
gcctctagaa agagctggga c 21

<210> 33
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:Synthetic
Oglionucleotide

<400> 33
cgccgtgttc catttatgag c 21

<210> 34
<211> 18
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:Synthetic
Oglionucleotide

<400> 34
gcatatttat gaatccca 18

<210> 35
<211> 18
<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
Oligonucleotide

<400> 35

actattaggg tcatgcac

18

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.